arXiv:1407.6232v1 [math.AP] 23 Jul 2014

EXISTENCE AND NONEXISTENCE OF LEAST ENERGY SOLUTIONS OF THE NEUMANN PROBLEM FOR A SEMILINEAR ELLIPTIC EQUATION WITH CRITICAL SOBOLEV EXPONENT AND A CRITICAL LOWER-ORDER PERTURBATION

DAVID G. COSTA AND PEDRO M. GIRÃO

Abstract. Let Ω be a smooth bounded domain in $\mathbb{R}^N$, with $N \geq 5$, $a > 0$, $\alpha \geq 0$ and $2^* = \frac{2N}{N-2}$. We show that the the exponent $q = \frac{2(N-1)}{N-2}$ plays a critical role regarding the existence of least energy (or ground state) solutions of the Neumann problem

$$\begin{cases} -\Delta u + au = u^{2^*-1} - \alpha u^{q-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases}$$

Namely, we prove that when $q = \frac{2(N-1)}{N-2}$ there exists an $\alpha_0 > 0$ such that the problem has a least energy solution if $\alpha < \alpha_0$ and has no least energy solution if $\alpha > \alpha_0$.

1. Introduction

Let Ω be a smooth bounded domain in $\mathbb{R}^N$, with $N \geq 5$, $a > 0$ and $\alpha \geq 0$. Let $2^* = \frac{2N}{N-2}$ be the critical exponent for the Sobolev embedding $H^1(\Omega) \subset L^q(\Omega)$ and $2^\# = \frac{2(N-1)}{N-2}$. We consider the problem

$$\begin{cases} -\Delta u + au = u^{2^*-1} - \alpha u^{q-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases} \qquad (\mathcal{P}_{\alpha,q})$$

We regard a as fixed and α as a parameter. From Theorem 3.2 of [18], due to X.J. Wang, we know that if $2 < q < 2^\#$, then problem $(\mathcal{P}_{\alpha,q})$ has a *least energy* solution for all values of $\alpha \geq 0$. (Wang's result actually holds for $N \geq 3$.) A question that naturally arises is the following: what happens for $q = 2^\#$?

It is well known that the solutions of $(\mathcal{P}_{\alpha,q})$ correspond to critical points of the functional $\Phi_\alpha : H^1(\Omega) \to \mathbb{R}$, defined by

$$\Phi_\alpha(u) := \frac{1}{2}|\nabla u|_2^2 + \frac{a}{2}|u|_2^2 + \frac{\alpha}{q}|u|_q^q - \frac{1}{2^*}|u|_{2^*}^{2^*},$$

2000 *Mathematics Subject Classification.* 46E35, 35J65.

Key words and phrases. Neumann problem, critical Sobolev exponent, least energy solutions, inequalities.

Pedro M. Girão was partially supported by FCT (Portugal).

where $|u|_p$ denotes the L^p norm of u in Ω. We recall that a *least energy* solution is a function $u \in H^1(\Omega)$ such that

$$\Phi_\alpha(u) = \inf_{\mathcal{N}} \Phi_\alpha.$$

The set $\mathcal{N}$ is the Nehari manifold, $\mathcal{N} := \{u \in H^1(\Omega) : \Phi'_\alpha(u)u = 0,\ u \neq 0\}$. It is interesting to note that when $q = 2^\#$ it is possible to determine explicitly the function $\Phi_\alpha|_{\mathcal{N}}$ by solving a quadratic equation. We take full advantage of this fact.

We recall that the infimum

$$S := \inf \left\{ \frac{\int_{\mathbb{R}^N} |\nabla u|^2}{(\int_{\mathbb{R}^N} |u|^{2^*})^{2/2^*}} \,\middle|\, u \in L^{2^*}(\mathbb{R}^N), \nabla u \in L^2(\mathbb{R}^N), u \neq 0 \right\}$$

is achieved by the Talenti instanton $U(x) := \left(\frac{N(N-2)}{N(N-2)+|x|^2}\right)^{\frac{N-2}{2}}$. For $\varepsilon > 0$ and $y \in \mathbb{R}^N$, we define $U_{\varepsilon,y} := \varepsilon^{-\frac{N-2}{2}} U\left(\frac{x-y}{\varepsilon}\right)$.

Heuristically, we can summarize the main idea behind the analysis of problem $(\mathcal{P}_{\alpha,q})$, when $q = 2^\#$, as follows. There exists an $\alpha_0 \in]0, +\infty]$ such that $\inf_{\mathcal{N}} \Phi_\alpha < \frac{S^{\frac{N}{2}}}{2N}$ for $\alpha < \alpha_0$, and $\inf_{\mathcal{N}} \Phi_\alpha = \frac{S^{\frac{N}{2}}}{2N}$ for $\alpha \geq \alpha_0$. If $\alpha < \alpha_0$, then $(\mathcal{P}_{\alpha,2^\#})$ has a least energy solution whereas, if $\alpha > \alpha_0$, then $(\mathcal{P}_{\alpha,2^\#})$ does not have a least energy solution. Suppose $\alpha_0 = +\infty$, so that there exist least energy solutions for all $\alpha \geq 0$. We choose a sequence $\alpha_k \to +\infty$ as $k \to +\infty$ and denote by u_k a corresponding sequence of least energy solutions. Then there would exist a sequence of positive numbers ε_k converging to zero, and a sequence of points $P_k \in \partial\Omega$, such that, modulo a subsequence, $P_k \to P$ and $|\nabla(u_k - U_{\varepsilon_k,P_k})|_2 \to 0$, as $k \to +\infty$. We can use $\Phi_{\alpha_k}(U_{\varepsilon_k,P_k})$ to estimate $\Phi_{\alpha_k}(u_k)$ from below with an error that is $o(\alpha_k\varepsilon_k)$. However, from Adimurthi and Mancini [1] and X.J. Wang [18], we have the estimate

$$\Phi_{\alpha_k}(U_{\varepsilon_k,P_k}) = \frac{S^{\frac{N}{2}}}{2N} - \frac{S^{\frac{N}{2}}}{2} H(P_k)A(N)\varepsilon_k + \frac{1}{2}B(N)\alpha_k\varepsilon_k + o(\alpha_k\varepsilon_k),$$

where $A(N)$ and $B(N)$ are positive constants that only depend on N, and $H(P_k)$ is the mean curvature of $\partial\Omega$ at P_k with respect to the unit outward normal. This lower bound is greater than $\frac{S^{\frac{N}{2}}}{2N}$, for large k. This contradicts the hypothesis that $\alpha_0 = +\infty$.

It is somewhat delicate to justify the use of $\Phi_{\alpha_k}(U_{\varepsilon_k,P_k})$ to estimate $\Phi_{\alpha_k}(u_k)$ from below. This was first done by Adimurthi, Pacella and Yadava in [2], who treated the case where $\alpha = 0$. The argument involves an expansion to second order of the energy at U_{ε_k,P_k} and a comparison of the eigenvalues of the linearized problem at U_{ε_k,P_k} with the eigenvalues of a limiting problem.

The present analysis builds on the work [2] of Adimurthi, Pacella and Yadava, which we will frequently refer to as [APY]. Of course, the works of Talenti [17], Brézis and Nirenberg [9], P.L. Lions [15], Adimurthi and Mancini [1], and X.J. Wang [18] are also of major importance for our study.

Our main result is the following

THEOREM. *Let Ω be a smooth bounded domain in $\mathbb{R}^N$, with $N \geq 5$, $a > 0$ and $\alpha \geq 0$. There exists a positive real number $\alpha_0 = \alpha_0(a, \Omega)$ such that*

(i) if $\alpha < \alpha_0$, then problem $(\mathcal{P}_{\alpha,2^\#})$ has a least energy solution;
(ii) if $\alpha > \alpha_0$, then problem $(\mathcal{P}_{\alpha,2^\#})$ has no least energy solution.

We remark that this result contrasts with Theorem 3.2 of [18], referred to above. Also, from this theorem we deduce an inequality (see (15)) which implies Aubin's inequality (16) (see [6] and Cherrier [11]).

We should mention that for any pair (a, α) (with $a > 0$ and $\alpha \geq 0$) problem $(\mathcal{P}_{\alpha,2^\#})$ has the constant solution

$$u = \kappa := \left(\tfrac{\alpha+\sqrt{\alpha^2+4a}}{2}\right)^{\frac{N-2}{2}}.$$

The energy of this solution is

$$\Phi_\alpha(\kappa) = \tfrac{|\Omega|}{2^\# N}\left[\left(\tfrac{\alpha+\sqrt{\alpha^2+4a}}{2}\right)^N + \tfrac{2^*}{2}a\left(\tfrac{\alpha+\sqrt{\alpha^2+4a}}{2}\right)^{N-2}\right],$$

where $|\Omega|$ denotes the N-dimensional Lebesgue measure of Ω. It follows that for $a > 0$ and $\alpha \geq 0$ sufficiently small, namely for $a \leq S/(2|\Omega|)^{\frac{2}{N}}$ and α such that $\Phi_\alpha(\kappa) \leq S^{\frac{N}{2}}/(2N)$, then the least energy solutions might be constant.

When the domain Ω is a ball and a is small, Adimurthi and Yadava [3] proved that $(\mathcal{P}_{0,2^\#})$ has more than one solution for $N = 4$, 5 and 6. However, when $N = 3$ a uniqueness result was proved by M. Zhu in [21] for convex domains, $\alpha = 0$ and small a.

Other works in the spirit of ours are those of Brézis and Lieb [8], Adimurthi and Yadava [4], M. Zhu [20], Z.Q. Wang [19] and Chabrowski and Willem [10].

The organization of this work is as follows. In Section 2 we give the setup of our work and the statement of the main result. In Section 3 we prove existence of least energy solutions. We then assume that the value α_0 is infinite and analyze the asymptotic behavior of the least energy solutions as $\alpha \to +\infty$. In Section 4 we prove nonexistence of least energy solutions. In Section 5 we give a lower bound for α_0 and, using the ideas of Chabrowski and Willem [10], give partial results concerning existence of least energy solutions for $\alpha = \alpha_0$. In Appendix A we check that the Nehari set $\mathcal{N}$ is a manifold and a natural constraint for Φ_α, we derive expressions for $\Phi_\alpha|_{\mathcal{N}}$, and we derive upper and lower bounds for $\Phi_\alpha|_{\mathcal{N}}$. Finally, in Appendix B we prove a technical estimate, used in our study, similar to those in Adimurthi and Mancini [1].

Motivated by this work, in [13] the second author has proved an inequality which improves inequality (15). In [14] he proves a family of inequalities which contains, as special cases, an inequality in Zhu's work [20] and the inequality in [13].

2. The setup and statement of the main result

Let Ω be a smooth bounded domain in $\mathbb{R}^N$, with $N \geq 5$. Let $2^* = \frac{2N}{N-2}$ be the critical exponent for the Sobolev embedding $H^1(\Omega) \subset L^q(\Omega)$ and $2^\# = \frac{2(N-1)}{N-2}$. Finally, let $a > 0$ and $\alpha \geq 0$. We are concerned with the

problem of existence of a *least energy* solution of

$$\begin{cases} -\Delta u + au = u^{2^*-1} - \alpha u^{2^\#-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases} \tag{1_α}$$

Solutions of (1_α) correspond to critical points of the functional $\Phi_\alpha : H^1(\Omega) \to \mathbb{R}$ defined by

$$\Phi_\alpha(u) := \frac{1}{2}||u||^2 + \frac{\alpha}{2^\#}|u|_{2^\#}^{2^\#} - \frac{1}{2^*}|u|_{2^*}^{2^*}. \tag{2}$$

We use the notations

$$|u|_p := (\int |u|^p)^{\frac{1}{p}} \qquad \text{and} \qquad ||u|| := \left(|\nabla u|_2^2 + a|u|_2^2\right)^{\frac{1}{2}}.$$

Unless otherwise indicated, integrals are over Ω.

We recall that the Nehari manifold is

$$\mathcal{N} := \left\{ u \in H^1(\Omega) : \Phi'_\alpha(u)u = 0, u \neq 0 \right\}.$$

For any $u \in H^1(\Omega) \setminus \{0\}$, there exists a unique $t(u) > 0$ such that $t(u)u \in \mathcal{N}$; the value of $t(u)$ is given in expression (62) of Appendix A. We define $\Psi_\alpha : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$ by

$$\Psi_\alpha(u) := \Phi_\alpha(t(u)u).$$

As can be checked in Appendix B,

$$\Psi_\alpha := \frac{1}{N}\frac{1}{2^\#}\frac{1}{2^N}\left[\left(\gamma + \sqrt{\gamma^2 + 4\beta}\right)^N + 2 \cdot 2^*\beta\left(\gamma + \sqrt{\gamma^2 + 4\beta}\right)^{N-2}\right], \tag{3}$$

where β, $\gamma : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$ are defined by

$$\beta(u) := \frac{||u||^2}{|u|_{2^*}^2} \tag{4}$$

and

$$\gamma(u) = \gamma_\alpha(u) := \alpha\frac{|u|_{2^\#}^{2^\#}}{|u|_{2^*}^{2^\#}}. \tag{5}$$

Equivalently,

$$\Psi_\alpha = \frac{1}{N}\frac{\beta^{\frac{N}{2}}}{2^\#}\left[\left(\delta + \sqrt{\delta^2 + 1}\right)^N + \frac{2^*}{2}\left(\delta + \sqrt{\delta^2 + 1}\right)^{N-2}\right], \tag{6}$$

with β as above and $\delta : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$ defined by

$$\delta(u) = \delta_\alpha(u) := \frac{\gamma(u)}{2\sqrt{\beta(u)}} = \frac{1}{2}\frac{\alpha|u|_{2^\#}^{2^\#}}{||u|| \cdot |u|_{2^*}^{\frac{2^*}{2}}}, \tag{7}$$

Obviously, every nonzero critical point of Φ_α is a critical point of Ψ_α. Since the Nehari manifold is a natural constraint for Φ_α, if u is a critical point of Ψ_α, then $t(u)u$ is a critical point of Φ_α.

As is usual, we say that $u \neq 0$ is a *ground state* critical point of Φ_α, or a *least energy* solution of (1_α), if

$$\Phi_\alpha(u) = \inf_{\mathcal{N}} \Phi_\alpha = \inf_{H^1(\Omega)\setminus\{0\}} \Psi_\alpha.$$

Our aim is to establish existence and nonexistence of least energy solutions of (1_α). We will consider the minimization problem corresponding to

$$S_\alpha := \inf \left\{ I_\alpha(u) | u \in H^1(\Omega) \setminus \{0\} \right\},$$

where $I_\alpha : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$ is defined by

$$I_\alpha := (N\Psi_\alpha)^{\frac{2}{N}}. \tag{8}$$

From (3) and (6) we obtain

$$I_\alpha = \frac{1}{4(2^\#)^{\frac{2}{N}}} \left[\left(\gamma + \sqrt{\gamma^2 + 4\beta} \right)^N + 2 \cdot 2^* \beta \left(\gamma + \sqrt{\gamma^2 + 4\beta} \right)^{N-2} \right]^{\frac{2}{N}} \tag{9}$$

and

$$I_\alpha = \frac{\beta}{(2^\#)^{\frac{2}{N}}} \left[\left(\delta + \sqrt{\delta^2 + 1} \right)^N + \frac{2^*}{2} \left(\delta + \sqrt{\delta^2 + 1} \right)^{N-2} \right]^{\frac{2}{N}}. \tag{10}$$

We observe that

$$I_\alpha \geq \beta, \tag{11}$$

since

$$\frac{1}{2^\#} \left(1 + \frac{2^*}{2} \right) = 1.$$

Before stating our main result, we recall that the infimum

$$S := \inf \left\{ \frac{\int_{\mathbb{R}^N} |\nabla u|^2}{(\int_{\mathbb{R}^N} |u|^{2^*})^{2/2^*}} \middle| u \in L^{2^*}(\mathbb{R}^N), \nabla u \in L^2(\mathbb{R}^N), u \neq 0 \right\},$$

which depends on N, is achieved by the Talenti instanton

$$U(x) := \left(\frac{N(N-2)}{N(N-2) + |x|^2} \right)^{\frac{N-2}{2}}.$$

This instanton U satisfies

$$-\Delta U = U^{2^*-1}, \tag{12}$$

so that

$$\int_{\mathbb{R}^N} |\nabla U|^2 = \int_{\mathbb{R}^N} U^{2^*} = S^{\frac{N}{2}}. \tag{13}$$

Let $\varepsilon > 0$ and $y \in \mathbb{R}^N$. For later use, we define the rescaled instanton

$$U_{\varepsilon,y} := \varepsilon^{-\frac{N-2}{2}} U \left(\frac{x-y}{\varepsilon} \right), \tag{14}$$

which also satisfies (12) and (13).

Our main result is

Theorem 2.1. *Let Ω be a smooth bounded domain in $\mathbb{R}^N$, with $N \geq 5$, $a > 0$ and $\alpha \geq 0$. There exists a positive real number $\alpha_0 = \alpha_0(a, \Omega)$ such that*

(i) if $\alpha < \alpha_0$, then (1_α) has a least energy solution;

(ii) if $\alpha > \alpha_0$, then (1_α) does not have a least energy solution and

$$\frac{S}{2^{\frac{2}{N}}} \leq \frac{\beta}{(2^\#)^{\frac{2}{N}}} \left[\left(\delta + \sqrt{\delta^2+1} \right)^N + \frac{2^*}{2} \left(\delta + \sqrt{\delta^2+1} \right)^{N-2} \right]^{\frac{2}{N}} \tag{15}$$

in $H^1(\Omega) \setminus \{0\}$, where β and $\delta = \delta_\alpha$ are defined in (4) and (7), respectively. The constant on the left hand side of (15) is sharp.

Corollary 2.2 (Aubin's inequality)**.** *Let Ω be a smooth bounded domain in $\mathbb{R}^N$, with $N \geq 5$. For every $\varsigma > 0$, there exists a $C(\varsigma, \Omega) > 0$ such that*

$$\frac{S}{2^{\frac{2}{N}}} - \varsigma \leq \frac{|\nabla u|_2^2 + C(\varsigma,\Omega)|u|_2^2}{|u|_{2^*}^2}, \tag{16}$$

for all $u \in H^1(\Omega) \setminus \{0\}$.

Proof. From Lemma 5.1, there exists a constant $\bar{c} > 0$ such that the right hand side of (15) is less than or equal to

$$\beta \left(1 + \frac{4}{2^\#}\delta + \bar{c}\delta^2 \right)$$

and from Hölder's inequality $|u|_{2^\#}^{2^\#} \leq |u|_2 |u|_{2^*}^{2^*/2}$. Hence $\delta(u) \leq \frac{\alpha}{2}\frac{|u|_2}{||u||}$. Let $\epsilon > 0$. For all $u \in H^1(\Omega)$,

$$\begin{aligned} \frac{S}{2^{\frac{2}{N}}}|u|_{2^*}^2 &\leq ||u||^2 \left(1 + \frac{2}{2^\#}\alpha_0 \frac{|u|_2}{||u||} + \bar{c}\frac{\alpha_0^2}{4}\frac{|u|_2^2}{||u||^2} \right) \\ &= ||u||^2 + \frac{2}{2^\#}\alpha_0 ||u||\,|u|_2 + \bar{c}\frac{\alpha_0^2}{4}|u|_2^2 \\ &\leq (1+\epsilon)|\nabla u|_2^2 + \left(\frac{\alpha_0^2}{(2^\#)^2\epsilon} + a\epsilon + \bar{c}\frac{\alpha_0^2}{4} \right) |u|_2^2. \end{aligned}$$

□

Remark 2.3. *Let $\kappa > 0$. By scaling, we easily check that*

$$\alpha_0 \left(\kappa^2 a, \frac{\Omega}{\kappa} \right) = \kappa\, \alpha_0(a, \Omega).$$

3. Existence of least energy solutions and their asymptotic behavior

In this section we start by proving the basic properties of the map $\alpha \mapsto S_\alpha$ and assertion (i) of Theorem 2.1. We then assume that the value α_0 in Theorem 2.1 is infinite and analyze the asymptotic behavior of the least energy solutions as $\alpha \to +\infty$.

As explained in the previous section, we consider the minimization problem corresponding to

$$S_\alpha := \inf\{I_\alpha(u) | u \in H^1(\Omega), u \neq 0\}.$$

From Adimurthi and Mancini [1] and X.J. Wang [18], we know that

$$0 < S_0 < \frac{S}{2^{\frac{2}{N}}} \tag{17}$$

(see (42) and (46) ahead). Obviously, S_α is nondecreasing as α increases. Choose any point $P \in \partial\Omega$. By testing I_α with $U_{\varepsilon,P}$ and letting $\varepsilon \to 0$, we conclude that $S_\alpha \leq \frac{S}{2^{\frac{2}{N}}}$ for all $\alpha \geq 0$.

Lemma 3.1. *If $S_\alpha < \frac{S}{2^{\frac{2}{N}}}$, then S_α is achieved.*

Proof. Let u_k be a minimizing sequence with $|u_k|_{2^*} = 1$. Since $\beta \leq I_\alpha$, from (11), (u_k) is bounded in $H^1(\Omega)$. We can assume that $u_k \rightharpoonup u$ in $H^1(\Omega)$, $u_k \to u$ a.e. on Ω, and $|\nabla(u_k - u)|^2 \rightharpoonup \mu$ and $|u_k - u|^{2^*} \rightharpoonup \nu$ in the sense of measures on $\bar{\Omega}$. Modulo a subsequence, the concentration-compactness lemma implies that

$$\lim_{k\to\infty} |\nabla u_k|_2^2 = |\nabla u|_2^2 + ||\mu||$$

and

$$\lim_{k\to\infty} |u_k|_{2^*}^{2^*} = |u|_{2^*}^{2^*} + ||\nu|| = 1,$$

where

$$\frac{S}{2^{\frac{2}{N}}} ||\nu||^{\frac{2}{2^*}} \leq ||\mu||.$$

This last inequality is an immediate consequence of inequality (16). For $S_\alpha = \lim_{k\to\infty} I_\alpha(u_k)$, we obtain that S_α equals

$$\frac{1}{4(2^\#)^{\frac{2}{N}}} \left[\left(\gamma_\infty + \sqrt{\gamma_\infty^2 + 4\beta_\infty} \right)^N + 2 \cdot 2^* \beta_\infty \left(\gamma_\infty + \sqrt{\gamma_\infty^2 + 4\beta_\infty} \right)^{N-2} \right]^{\frac{2}{N}},$$

with

$$\beta_\infty = ||u||^2 + ||\mu|| = \frac{||u||^2 + ||\mu||}{(|u|_{2^*}^{2^*} + ||\nu||)^{\frac{2}{2^*}}}$$

and

$$\gamma_\infty = \alpha |u|_{2^\#}^{2^\#} = \alpha \frac{|u|_{2^\#}^{2^\#}}{(|u|_{2^*}^{2^*} + ||\nu||)^{\frac{2^\#}{2^*}}}.$$

If $u = 0$, then

$$\beta_\infty = \frac{||\mu||}{||\nu||^{\frac{2}{2^*}}} \geq \frac{S}{2^{\frac{2}{N}}},$$

a contradiction. So $u \neq 0$.

We claim that $||\mu|| = 0$. We argue by contradiction and suppose that $||\mu|| \neq 0$. If $||\nu|| = 0$, then $S_\alpha > I_\alpha(u)$, which is impossible. So $||\nu|| \neq 0$.

Let $x_0 := |u|_{2^*}^{2^*}$, so that $1 - x_0 = ||\nu||$. We define f, g and $h : [0,1] \to \mathbb{R}$ by

$$f(x) := \underline{\gamma} x^{\frac{2^\#}{2^*}} + \sqrt{\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}} + 4\underline{\beta} x^{\frac{2}{2^*}} + 4\frac{||\mu||}{||\nu||^{2/2^*}}(1-x)^{\frac{2}{2^*}}},$$

$$g(x) := \underline{\beta} x^{\frac{2}{2^*}} + \frac{||\mu||}{||\nu||^{2/2^*}}(1-x)^{\frac{2}{2^*}}$$

and

$$h := f^N + 2 \cdot 2^* f^{N-2} g,$$

for $\underline{\beta} = \beta(u)$ and $\underline{\gamma} = \gamma(u)$. The value S_α is

$$S_\alpha = \frac{1}{4(2^\#)^{\frac{2}{N}}} \left[h(x_0) \right]^{\frac{2}{N}}.$$

We wish to prove that the minimum of h occurs at 0 or 1. The former case corresponds to $u = 0$ and the latter to $||\nu|| = 0$. In either case we are led to a contradiction. This will prove that $||\mu|| = 0$, thereby establishing the claim.

The derivative of h is

$$h' = f^{N-3}[N(f^2 + 4g)f' + 2 \cdot 2^* f g'].$$

Since

$$f^2 + 4g = 2f\sqrt{\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}} + 4g},$$

we can write

$$h' = 2f^{N-2}\left[N\sqrt{\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}} + 4g}\, f' + 2^* g'\right].$$

The expression for h' can be further simplified by computing f':

$$\begin{aligned}\sqrt{\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}} + 4g}\, f' &= \frac{1}{2^*}\left[2^\#\underline{\gamma} x^{\frac{2^\#}{2^*}-1}\sqrt{\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}} + 4g} + 2^\#\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}-1} + 4g'\right] \\ &= \frac{2^\#}{2^*}\left[\underline{\gamma} x^{\frac{2^\#}{2^*}-1} f + 2\frac{2^*}{2^\#} g'\right].\end{aligned}$$

This yields

$$\begin{aligned}h' &= 2(N-1)f^{N-2}\left[\underline{\gamma} x^{\frac{2^\#}{2^*}-1} f + 2^* g'\right] \\ &= 2(N-1)f^{N-2} x^{\frac{2^\#}{2^*}-1}\left[\underline{\gamma} f + 2^* x^{1-\frac{2^\#}{2^*}} g'\right].\end{aligned}$$

We notice that $h'(0) = +\infty$ and $h'(1) = -\infty$; at a zero of h', $g' < 0$.

At a point of minimum of h in the interior of $[0, 1]$, $h' = 0$ and

$$\begin{aligned}\sqrt{\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}} + 4g}\, f' &= \frac{2^\#}{2^*}\left[\underline{\gamma} x^{\frac{2^\#}{2^*}-1} f + 2^* g' - 2^*\left(1 - \frac{2}{2^\#}\right) g'\right] \\ &= -(2^\# - 2) g';\end{aligned}$$

we notice that at a zero of h', $f' > 0$.

We consider

$$\kappa := -2^* x^{1-\frac{2^\#}{2^*}} g',$$

whose derivative is

$$\begin{aligned}\kappa' &= -(2^* - 2^\#) x^{-\frac{2^\#}{2^*}} g' - 2^* x^{\frac{2^*-2^\#}{2^*}} g'' \\ &> -(2^* - 2^\#) x^{-\frac{2^\#}{2^*}} g' \\ &= x^{-\frac{2^\#}{2^*}}\sqrt{\underline{\gamma}^2 x^{2\frac{2^\#}{2^*}} + 4g}\, f' \quad \text{for } h' = 0 \\ &> \underline{\gamma} f'.\end{aligned}$$

The zeros of h' occur when $\underline{\gamma} f = \kappa$. We just proved that $\kappa' > \underline{\gamma} f'$ at the zeros of h'. This implies that the graphs of $\underline{\gamma} f$ and κ can cross at most once, and that h' has at most one zero. If the function h were to have a minimum in the interior of $[0, 1]$, then h' would have at least three zeros because $h'(0) = +\infty$ and $h'(1) = -\infty$. We conclude that h has no minimum

inside $[0,1]$. (The conditions on the derivative of h at the end points of the interval, or the fact that the graphs of $\underline{\gamma} f$ and κ cross, imply that h' does vanish inside $[0,1]$, at a point of maximum of h.) Therefore the minimum of h occurs either at 0 or 1 and we have proved our claim.

Since $||\mu|| = 0$, the function u is a minimizer for I_α. □

Lemma 3.2. *The map $\alpha \mapsto S_\alpha$ is continuous on $[0, +\infty[$.*

Proof. Let $\bar{\alpha} \in [0, +\infty[$. First we prove that $\alpha \mapsto S_\alpha$ is continuous from the right at $\bar{\alpha}$. If $S_{\bar{\alpha}} = \frac{S}{2^{\frac{2}{N}}}$, then continuity from the right at $\bar{\alpha}$ is obvious. If $S_{\bar{\alpha}} < \frac{S}{2^{\frac{2}{N}}}$, let $u_{\bar{\alpha}}$ be a minimizer of $I_{\bar{\alpha}}$, which exists by the previous lemma. If $\alpha > \bar{\alpha}$, then $S_{\bar{\alpha}} \leq S_\alpha \leq I_\alpha(u_{\bar{\alpha}}) \to S_{\bar{\alpha}}$ as $\alpha \searrow \bar{\alpha}$. This proves continuity from the right at $\bar{\alpha}$.

To prove continuity from the left we show that $\lim_{\alpha \nearrow \bar{\alpha}} S_\alpha = S_{\bar{\alpha}}$. If the value of the limit on the left hand side is $\frac{S}{2^{\frac{2}{N}}}$, then this equality is obvious. So suppose $\lim_{\alpha \nearrow \bar{\alpha}} S_\alpha < \frac{S}{2^{\frac{2}{N}}}$. Choose a sequence $\alpha_k \nearrow \bar{\alpha}$ and $u_k \in H^1(\Omega)$, with $|u_k|_{2^*} = 1$, such that $I_{\alpha_k}(u_k) = S_{\alpha_k}$. By (11), the sequence (u_k) is bounded in $H^1(\Omega)$ and we can assume that $u_k \rightharpoonup u$ in $H^1(\Omega)$. An application of the concentration-compactness principle, as in the previous lemma, shows that $u \neq 0$ and

$$\lim_{k\to\infty} I_{\alpha_k}(u_k) \geq I_{\bar{\alpha}}(u).$$

So,

$$S_{\bar{\alpha}} \leq I_{\bar{\alpha}}(u) \leq \lim_{k\to\infty} I_{\alpha_k}(u_k) = \lim_{\alpha_k \nearrow \bar{\alpha}} S_\alpha$$

and $S_{\bar{\alpha}} = \lim_{\alpha_k \nearrow \bar{\alpha}} S_\alpha$. □

By the previous lemma, the value

$$\alpha_0 := \begin{cases} +\infty, \text{ if } S_\alpha < \frac{S}{2^{\frac{2}{N}}} \text{ for all } \alpha \in [0, +\infty[, \\ \min\left\{\alpha \in [0, +\infty[\;\middle|\; S_\alpha = \frac{S}{2^{\frac{2}{N}}}\right\}, \text{ otherwise.} \end{cases} \tag{18}$$

is well defined. By (17) it is not zero. Lemma 3.1 implies the following two corollaries:

Corollary 3.3. *The map $\alpha \mapsto S_\alpha$ is strictly increasing on $[0, \alpha_0]$.*

Corollary 3.4. *If $\alpha \in [0, \alpha_0[$, then (1_α) has a least energy solution u_α. If $\alpha \in]\alpha_0, +\infty[$, then (1_α) does not have a least energy solution.*

This proves (i) of Theorem 2.1. Assertion (ii) of Theorem 2.1 will also follow once we establish that α_0 is finite.

Lemma 3.5. *If $S_\alpha < \frac{S}{2^{\frac{2}{N}}}$ for all $\alpha \geq 0$, then*

$$\lim_{\alpha\to+\infty} S_\alpha = \frac{S}{2^{\frac{2}{N}}}. \tag{19}$$

Suppose $\alpha_k \to +\infty$ as $k \to +\infty$ and u_k is a minimizer for I_{α_k} satisfying (1_{α_k}). Then $u_k \rightharpoonup 0$ in $H^1(\Omega)$ and

$$M_k := \max_{\bar{\Omega}} u_k$$

converges to $+\infty$, as $k \to \infty$.

Proof. Suppose $S_\alpha < \frac{S}{2^{\frac{2}{N}}}$ for all $\alpha \geq 0$ and choose any sequence $\alpha_k \to +\infty$ as $k \to +\infty$. Let u_k be a minimizer for I_{α_k} satisfying (1_{α_k}), which necessarily exists by Lemma 3.1 and rescaling. We claim that u_k is bounded in $H^1(\Omega)$. Indeed, by (9),

$$\frac{1}{(2^\#)^{\frac{2}{N}}}\,\gamma^2(u_k) \leq I_\alpha(u_k) \leq \frac{S}{2^{\frac{2}{N}}}.$$

So,

$$\alpha_k |u_k|_{2^\#}^{2^\#} \leq \left(\frac{2^\#}{2}\right)^{\frac{1}{N}} S^{\frac{1}{2}} |u_k|_{2^*}^{2^\#}.$$

By (1_{α_k}),

$$|u_k|_{2^*}^{2^*} = ||u_k||^2 + \alpha_k |u_k|_{2^\#}^{2^\#}. \tag{20}$$

Together,

$$\begin{aligned} |u_k|_{2^*}^{2^*-2} &\leq \beta(u_k) + \left(\frac{2^\#}{2}\right)^{\frac{1}{N}} S^{\frac{1}{2}} |u_k|_{2^*}^{2^\#-2} \\ &\leq \frac{S}{2^{\frac{2}{N}}} + \left(\frac{2^\#}{2}\right)^{\frac{1}{N}} S^{\frac{1}{2}} |u_k|_{2^*}^{2^\#-2}, \end{aligned}$$

since, by (11), $\beta(u_k) \leq I_{\alpha_k}(u_k) \leq \frac{S}{2^{\frac{2}{N}}}$. So $|u_k|_{2^*}$ is bounded. Recalling that $\beta(u_k) = \frac{||u_k||^2}{|u_k|_{2^*}^2}$, we conclude that u_k is bounded in $H^1(\Omega)$.

From (20), we conclude that $u_k \rightharpoonup 0$ in $H^1(\Omega)$. We can assume that $u_k \to 0$ a.e. on Ω, and $|\nabla u_k|^2 \rightharpoonup \mu$ and $|u_k|^{2^*} \rightharpoonup \nu$ in the sense of measures on $\bar{\Omega}$. Then

$$\lim_{k\to\infty} |\nabla u_k|_2^2 = ||\mu|| \tag{21}$$

and

$$\lim_{k\to\infty} |u_k|_{2^*}^{2^*} = ||\nu||, \tag{22}$$

where

$$\frac{S}{2^{\frac{2}{N}}} ||\nu||^{\frac{2}{2^*}} \leq ||\mu||. \tag{23}$$

Thus

$$\frac{S}{2^{\frac{2}{N}}} \geq \lim_{k\to\infty} S_{\alpha_k} = \lim_{k\to\infty} I_{\alpha_k}(u_k) \geq \frac{||\mu||}{||\nu||^{\frac{2}{2^*}}} \geq \frac{S}{2^{\frac{2}{N}}}, \tag{24}$$

and the inequalities in (24) are equalities. This proves (19).

From (1_{α_k}), the values M_k satisfy

$$a + \alpha_k M_k^{2^\#-2} \leq M_k^{2^*-2} \tag{25}$$

and consequently $M_k \to +\infty$ as $k \to +\infty$. □

Lemma 3.6. *Let $S_{\alpha_k} < \frac{S}{2^{\frac{2}{N}}}$ and $S_{\alpha_k} \to \frac{S}{2^{\frac{2}{N}}}$ as $\alpha_k \to \alpha_0 \in]0, +\infty]$. Denote by $u_k \in H^1(\Omega)$ a minimizer for I_{α_k} satisfying (1_{α_k}). In case $\alpha_0 < +\infty$ suppose that $u_k \rightharpoonup 0$. Then*

$$\lim_{k\to\infty} |\nabla u_k|_2^2 = \lim_{k\to\infty} |u_k|_{2^*}^{2^*} = \frac{S^{\frac{N}{2}}}{2}. \tag{26}$$

Moreover, if $\alpha_0 = +\infty$, *or if* $\alpha_0 < +\infty$ *and we further assume that* $\lim_{\alpha_k \to \alpha_0} M_k = +\infty$, *then we also have*

$$\lim_{k\to\infty} \alpha_k \delta_k = 0, \tag{27}$$

$$\lim_{k\to\infty} |\nabla u_k - \nabla U_{\delta_k, P_k}|_2 = 0 \tag{28}$$

and $P_k \in \partial\Omega$, *for large* k. *Here, we are denoting*

$$\delta_k := M_k^{-\frac{2}{N-2}},$$

and P_k *is such that* $M_k = u_k(P_k)$.

Note. If $\alpha_0 = +\infty$, Lemma 3.5 guarantees the conditions $S_{\alpha_k} \to \frac{S}{2^{\frac{2}{N}}}$, $u_k \rightharpoonup 0$ and $M_k \to +\infty$ are satisfied.

Proof. By (1_{α_k}), u_k satisfies (20). Since $u_k \rightharpoonup 0$ in $H^1(\Omega)$, u_k is bounded in $H^1(\Omega)$. Therefore, (21), (22), (23) and (24) hold, with equalities in (24). Hence, $\beta(u_k) \to \frac{S}{2^{\frac{2}{N}}}$. From (10), $\delta(u_k) \to 0$ and

$$\lim_{k\to\infty} \alpha_k |u_k|_{2^\#}^{2^\#} = 0,$$

as u_k is bounded in $H^1(\Omega)$. Taking limits in (20) as $k \to \infty$,

$$||\nu|| = ||\mu||. \tag{29}$$

Combining (24) and (29), equalities (26) follow.

We now use the Gidas and Spruck blow up technique [12]. Let $v_k(x) := \delta_k^{\frac{N-2}{2}} u_k(\delta_k x + P_k)$ for $x \in \Omega_k := (\Omega - P_k)/\delta_k$, so that

$$\begin{cases} -\Delta v_k + a\delta_k^2 v_k + \alpha_k \delta_k v_k^{2^\#-1} = v_k^{2^*-1} & \text{in } \Omega_k, \\ 0 < v_k \le v_k(0) = 1 & \text{in } \Omega_k, \\ \frac{\partial v_k}{\partial \nu} = 0 & \text{on } \partial\Omega_k. \end{cases}$$

Rewriting (25) in terms of the δ_k,

$$a\delta_k^2 + \alpha_k \delta_k \le 1.$$

So, we can assume that $P_k \to P_0$,

$$\operatorname{dist}(P_k, \partial\Omega)/\delta_k \to L \in [0, +\infty],$$

$$\Omega_k \to \Omega_\infty := \{(\tilde{x}, x_N) \in \mathbb{R}^{N-1} \times \mathbb{R} : x_N > -L\}$$

and $\alpha_k \delta_k \to \bar{\alpha}$. By the elliptic estimates in [5],

$$v_k \to v \text{ in } C^2_{\text{loc}}(\Omega_\infty) \tag{30}$$

where v satisfies

$$\begin{cases} -\Delta v + \bar{\alpha} v^{2^\#-1} = v^{2^*-1} & \text{in } \Omega_\infty, \\ 0 < v \le v(0) = 1 & \text{in } \Omega_\infty, \\ \frac{\partial v}{\partial \nu} = 0 & \text{on } \partial\Omega_\infty \end{cases}$$

as $a\delta_k^2 \to 0$. By lower semicontinuity of the norm, $v \in L^{2^*}(\Omega_\infty)$ and $\nabla v \in L^2(\Omega_\infty)$. So, we can apply Pohozaev's identity and get $\bar{\alpha} = 0$, and thus $v = U$.

If $L = +\infty$, then $\Omega_\infty = \mathbb{R}^N$. From (26),

$$S^{\frac{N}{2}} = \int_{\mathbb{R}^N} |\nabla U|^2 \leq \lim_{k\to\infty} |\nabla u_k|_2^2 = \frac{S^{\frac{N}{2}}}{2},$$

which is impossible.

So L is finite. This implies that $P_0 \in \partial\Omega$. In fact, L has to be zero since $v \leq v(0)$. Using a diffeomorphism to straighten a boundary portion of Ω, the argument in Lemma 2.2 of [APY] shows that $P_k \in \partial\Omega$ for large k. Finally, from (26), (30) and

$$\int_{\mathbb{R}^N_+} |\nabla U|^2 = \frac{S^{\frac{N}{2}}}{2},$$

we deduce (28). □

As in [2] and [7], let

$$\mathcal{M} := \{CU_{\varepsilon,y}, C \in \mathbb{R}, \varepsilon > 0, y \in \partial\Omega\}$$

and $d(u, \mathcal{M}) := \inf\{|\nabla(u - V)|_2, V \in \mathcal{M}\}$. The set $\mathcal{M} \setminus \{0\}$ is a manifold of dimension $N + 1$. The tangent space $T_{C_l,\varepsilon_l,y_l}(\mathcal{M})$ at $C_l U_{\varepsilon_l,y_l}$ is given by

$$T_{C_l,\varepsilon_l,y_l}(\mathcal{M}) = \text{span}\left\{U_{\varepsilon,y}, C\frac{\partial}{\partial\varepsilon}U_{\varepsilon,y}, C\frac{\partial}{\partial\tau_i}U_{\varepsilon,y}, 1 \leq i \leq N-1\right\}_{(C_l,\varepsilon_l,y_l)}$$

where $T_x(\partial\Omega) = \text{span}\{\tau_1, \ldots, \tau_{N-1}\}$.

For large k, the infimum $d(u_k, \mathcal{M})$ is achieved:

$$d(u_k, \mathcal{M}) = |\nabla(u_k - C_k U_{\varepsilon_k,y_k})|_2 \text{ for } C_k U_{\varepsilon_k,y_k} \in \mathcal{M}. \tag{31}$$

Furthermore,

$$C_k = 1 + o(1) \tag{32}$$

$y_k \to P_0$ and $\varepsilon_k/\delta_k \to 1$ (see Lemma 1 of [7] and Lemma 2.3 of [2]). From (27),

$$\alpha_k \varepsilon_k \to 0. \tag{33}$$

We define

$$w_k := u_k - C_k U_{\varepsilon_k,y_k},$$

so that

$$\int \nabla U_{\varepsilon_k,y_k} \cdot \nabla w_k = 0. \tag{34}$$

Now, on the one hand, from (28),

$$\lim_{k\to\infty} |\nabla(u_k - C_k U_{\varepsilon_k,y_k})|_2 = 0.$$

On the other hand, from Poincaré's inequality, and the fact that both the average of u_k and the average of $C_k U_{\varepsilon_k,y_k}$, in Ω, converge to zero,

$$\lim_{k\to\infty} |u_k - C_k U_{\varepsilon_k,y_k}|_{2^*} = 0.$$

Together,

$$\lim_{k\to\infty} ||w_k|| = 0. \tag{35}$$

Our next objective is the upper bound in Lemma 3.11 for $\int U_{\varepsilon_k,y_k}^{2^*-2} w_k^2$ in terms of $|\nabla w_k|_2^2 + (2^\# - 1)\alpha_k \int U_{\varepsilon_k,y_k}^{2^\#-2} w_k^2$. This will be crucial in the lower bound estimates for the energy in Section 4.

The eigenvalue problems arising from the linearization of (1_{α_k}) at U_{ε_k,y_k} are related to the eigenvalue problem in

Lemma 3.7 (Bianchi and Egnell [7], Rey [16]). *The eigenvalue problem*

$$\begin{cases} -\Delta\varphi = \mu U^{2^*-2}\varphi & \text{in } \mathbb{R}^N_+, \\ \frac{\partial\varphi}{\partial\nu} = 0 & \text{on } \partial\mathbb{R}^N_+, \\ \int_{\mathbb{R}^N_+} U^{2^*-2}\varphi^2 < \infty \end{cases}$$

admits a discrete spectrum $\mu_1 < \mu_2 \le \mu_3 \le \dots$ such that $\mu_1 = 1$, $\mu_2 = \mu_3 = \dots = \mu_N = 2^* - 1$ and $\mu_{N+1} > 2^* - 1$. The eigenspaces V_1 and $V_{(2^*-1)}$, corresponding to 1 and $(2^* - 1)$, are given by

$$\begin{aligned} V_1 &= \text{span } U, \\ V_{(2^*-1)} &= \text{span}\left\{ \frac{\partial U_{1,y}}{\partial y_i}\Big|_{y=0}, \text{ for } 1 \le i \le N-1 \right\}. \end{aligned}$$

We will consider the eigenvalue problems arising from the linearization of (1_{α_k}) at U_{ε_k,y_k}. Let $\varepsilon > 0$, $\nu_\varepsilon > 0$, and $y_\varepsilon \in \partial\Omega$ with $\lim_{\varepsilon\to 0} y_\varepsilon = y_0$. Let $\{\varphi_{i,\varepsilon}\}_{i=1}^\infty$ be a complete set of orthogonal eigenfunctions with eigenvalues $\mu_{1,\varepsilon} < \mu_{2,\varepsilon} \le \mu_{3,\varepsilon} \le \dots$ for the weighted eigenvalue problem

$$\begin{cases} -\Delta\varphi + \nu_\varepsilon U^{2^\#-2}_{\varepsilon,y_\varepsilon}\varphi = \mu U^{2^*-2}_{\varepsilon,y_\varepsilon}\varphi & \text{in } \Omega, \\ \frac{\partial\varphi}{\partial\nu} = 0 & \text{on } \partial\Omega, \end{cases}$$

with $\varphi_{1,\varepsilon} > 0$ and

$$\int_\Omega U^{2^*-2}\varphi_{i,\varepsilon}\varphi_{j,\varepsilon} = \delta_{i,j}.$$

Let

$$\Omega_\varepsilon := (\Omega - y_\varepsilon)/\varepsilon.$$

The sets Ω_ε converge to a half space as $\varepsilon \to 0$. For a function v on Ω, we define $\tilde v$ on Ω_ε by

$$\tilde v(x) := \varepsilon^{\frac{N-2}{2}} v(\varepsilon x + y_\varepsilon).$$

The relation between these eigenvalue problems and the one considered in Lemma 3.7 is given in

Lemma 3.8. *Suppose $y_\varepsilon \in \partial\Omega$, $\lim_{\varepsilon\to 0} y_\varepsilon = y_0$, $\lim_{\varepsilon\to 0}(\varepsilon\nu_\varepsilon) = 0$ and the sets Ω_ε converge to $\mathbb{R}^N_+$. Then, up to a subsequence,*

$$\lim_{\varepsilon\to 0} \mu_{i,\varepsilon} = \mu_i$$

and

$$\lim_{\varepsilon\to 0} \int_{\Omega_\varepsilon} U^{2^*-2}(\tilde\varphi_{i,\varepsilon} - \tilde\varphi_i)^2 = 0.$$

The μ_i and $\tilde\varphi_i$ satisfy

$$\begin{cases} -\Delta\tilde\varphi_i = \mu_i U^{2^*-2}\tilde\varphi_i & \text{in } \mathbb{R}^N_+, \\ \frac{\partial\tilde\varphi_i}{\partial\nu} = 0 & \text{on } \partial\mathbb{R}^N_+, \\ \int_{\mathbb{R}^N_+} U^{2^*-2}\tilde\varphi_i^2 = 1, \end{cases}$$

and the functions $\tilde\varphi_i$ are supposed extended to $\mathbb{R}^N$ by reflection. In particular, from the previous lemma, $\mu_1 = 1$, $\tilde\varphi_1 = CU$ for some constant $C > 0$, $\mu_i = 2^* - 1$ for $2 \le i \le N$ and $\mu_{N+1} > 2^* - 1$. Also, $\{\tilde\varphi_i\}_{i=2}^N$ is in the span of $\{\partial U_{1,y}/\partial y_i|_{y=0}, \text{ for } 1 \le i \le N-1\}$.

The proof of Lemma 3.8 is a consequence of the arguments in the proof of Lemma 3.3 of [APY], of Lemma 3.9 and of Remark 3.10. For the details we refer to the proof of Lemma 5.5 of [14] for parameter s there equal to one.

Lemma 3.9. *Suppose* $y_\varepsilon \in \bar\Omega$, $\varphi_\varepsilon \in H^1(\Omega)$,

$$\begin{cases} \displaystyle\int U_{\varepsilon,y_\varepsilon}^{2^\#-2}\varphi_\varepsilon^2 \to 0, \\ \\ \displaystyle\int |\nabla\varphi_\varepsilon|^2 \to 0, \end{cases}$$

as $\varepsilon \to 0$. *Then*

$$\int U_{\varepsilon,y_\varepsilon}^{2^*-2}\varphi_\varepsilon^2 \to 0,$$

as $\varepsilon \to 0$.

Proof. We denote the average of φ_ε in Ω by $\bar\varphi_\varepsilon$. By Poincaré's inequality,

$$|\varphi_\varepsilon - \bar\varphi_\varepsilon|_{2^*} \to 0.$$

The limits in this proof are taken as $\varepsilon \to 0$. So we can write $\varphi_\varepsilon = \bar\varphi_\varepsilon + \eta_\varepsilon$, with $\eta_\varepsilon \to 0$ in L^{2^*}. We know that

$$\int U_{\varepsilon,y_\varepsilon}^{2^\#-2}(\bar\varphi_\varepsilon^2 + 2\bar\varphi_\varepsilon\eta_\varepsilon + \eta_\varepsilon^2) = o(1).$$

We have the following estimates for the three terms on the left hand side:

$$\int U_{\varepsilon,y_\varepsilon}^{2^\#-2}\bar\varphi_\varepsilon^2 \ge b\bar\varphi_\varepsilon^2\varepsilon,$$

for some $b > 0$, and

$$\begin{aligned} \left|\int U_{\varepsilon,y_\varepsilon}^{2^\#-2}\eta_\varepsilon\bar\varphi_\varepsilon\right| &\le |\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\left(\int U_{\varepsilon,y_\varepsilon}^{\frac{N}{N-2}\frac{4}{N+2}}\right)^{\frac{N+2}{2N}} \\ &\le C|\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\varepsilon, \end{aligned}$$

by (39); and

$$\begin{aligned} \int U_{\varepsilon,y_\varepsilon}^{2^\#-2}\eta_\varepsilon^2 &\le |\eta_\varepsilon|_{2^*}^2\left(\int U_{\varepsilon,y_\varepsilon}^{\frac{N}{N-2}}\right)^{\frac{2}{N}} \\ &\le C|\eta_\varepsilon|_{2^*}^2\varepsilon|\log\varepsilon|^{\frac{2}{N}}, \end{aligned} \tag{36}$$

by (40). (Inequalities (39), (40) and (41) are in the beginning of the next section.) Thus

$$b\bar\varphi_\varepsilon^2\varepsilon \le C|\bar\varphi_\varepsilon|\varepsilon + o(1).$$

This shows that $\bar\varphi_\varepsilon\sqrt{\varepsilon}$ is bounded. But if $\bar\varphi_\varepsilon\sqrt{\varepsilon}$ is bounded this shows that

$$\bar\varphi_\varepsilon\sqrt{\varepsilon} \to 0. \tag{37}$$

We want to prove that

$$\int U_{\varepsilon,y_\varepsilon}^{2^*-2}(\bar\varphi_\varepsilon^2 + 2\bar\varphi_\varepsilon\eta_\varepsilon + \eta_\varepsilon^2) = o(1).$$

For the first term on the left hand side we have, by (39) and then (37),

$$\int U_{\varepsilon,y_\varepsilon}^{2^*-2}\bar\varphi_\varepsilon^2 \le C\bar\varphi_\varepsilon^2\varepsilon^2 \to 0.$$

For the third term we have

$$\int U_{\varepsilon,y_\varepsilon}^{2^*-2}\eta_\varepsilon^2 \le C|\eta_\varepsilon|_{2^*}^2 \to 0.$$

We claim that the remaining term also converges to zero. This will prove the lemma. For the second term we have the estimate

$$\zeta_\varepsilon := \left|\int U_{\varepsilon,y_\varepsilon}^{2^*-2}\bar\varphi_\varepsilon\eta_\varepsilon\right| \le |\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\left(\int U_{\varepsilon,y_\varepsilon}^{\frac{N}{N-2}\frac{8}{N+2}}\right)^{\frac{N+2}{2N}}.$$

If $N=5$, by (41),

$$\zeta_\varepsilon \le C|\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\varepsilon^{N\left(1-\frac{4}{N+2}\right)\frac{N+2}{2N}} \le C|\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\varepsilon^{\frac{N-2}{2}} = C|\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\varepsilon^{\frac{3}{2}}.$$

If $N=6$, by (40),

$$\zeta_\varepsilon \le C|\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\varepsilon^2|\log\varepsilon|^{\frac{2}{3}}.$$

Finally, if $N\ge 7$, by (39),

$$\zeta_\varepsilon \le C|\eta_\varepsilon|_{2^*}|\bar\varphi_\varepsilon|\varepsilon^2.$$

In all three cases, (37) implies that $\zeta_\varepsilon\to 0$. □

Remark 3.10. *If in the previous lemma, instead of assuming $\int|\nabla\varphi_\varepsilon|^2\to 0$, we assume that $\int|\nabla\varphi_\varepsilon|^2$ is bounded, then we can still conclude $\bar\varphi_\varepsilon\sqrt{\varepsilon}\to 0$ and $\int U_{\varepsilon,y_\varepsilon}^{2^*-2}\left(\bar\varphi_\varepsilon^2+2\bar\varphi_\varepsilon\eta_\varepsilon\right) = \int_{\Omega_\varepsilon}U^{2^*-2}\left(\tilde{\bar\varphi}_\varepsilon^2+2\tilde{\bar\varphi}_\varepsilon\tilde\eta_\varepsilon\right)\to 0$, as $\varepsilon\to 0$.*

Using Lemma 3.8 and the arguments in the proof of Lemma 3.4 of [APY], we deduce

Lemma 3.11. *Suppose $y_\varepsilon\in\partial\Omega$, $\lim_{\varepsilon\to 0}y_\varepsilon = y_0$ and $\lim_{\varepsilon\to 0}(\varepsilon\nu_\varepsilon)=0$. There exists a constant $\gamma_1>0$ such that, for sufficiently small ε,*

$$|\nabla w|_2^2+\nu_\varepsilon\int U_{\varepsilon,y_\varepsilon}^{2^\#-2}w^2 \ge (2^*-1+\gamma_1)\int U_{\varepsilon,y_\varepsilon}^{2^*-2}w^2 + O(\varepsilon^2||w||^2)$$

for w orthogonal to $T_{1,\varepsilon,y_\varepsilon}(\mathcal{M})$.

4. Nonexistence of least energy solutions

In this section we prove (ii) of Theorem 2.1. The idea of the proof is to obtain a lower bound for I_α and show that if α_0, defined in (18), is infinite, then the least energy solutions u_k of (1_{α_k}) have energy $I_{\alpha_k}(u_k)>\frac{S}{2^{\frac{2}{N}}}$, for large α_k. This is impossible. Therefore α_0 is finite. By Corollary 3.4, (ii) of Theorem 2.1 follows.

Assume

$$u_k = C_kU_{\varepsilon_k,y_k}+w_k,$$

(26), (31), (32), (33) and (35). From (10), I_α has the lower bound

$$I_\alpha \ge \beta\left(1+\frac{4}{2^\#}\delta\right) \tag{38}$$

(this is also checked in (63) of Appendix A). We will expand β and δ to second order around U_{ε_k,y_k}. We start by deriving estimates for the terms that appear in this expansion.

We recall, from Brézis and Nirenberg [9], that, for $y \in \bar{\Omega}$, there exist positive constants c_1 and c_2 such that:
if $1 \leq q < \frac{N}{N-2}$, then

$$c_1 \varepsilon^{q\left(\frac{N-2}{2}\right)} \leq |U_{\varepsilon,y}|_q^q \leq c_2 \varepsilon^{q\left(\frac{N-2}{2}\right)}; \tag{39}$$

if $q = \frac{N}{N-2}$, then

$$c_1 \varepsilon^{\frac{N}{2}} |\log \varepsilon| \leq |U_{\varepsilon,y}|_q^q \leq c_2 \varepsilon^{\frac{N}{2}} |\log \varepsilon|; \tag{40}$$

and if $\frac{N}{N-2} < q \leq \frac{2N}{N-2}$, then

$$c_1 \varepsilon^{N\left(1-\frac{q}{2^*}\right)} \leq |U_{\varepsilon,y}|_q^q \leq c_2 \varepsilon^{N\left(1-\frac{q}{2^*}\right)}. \tag{41}$$

For brevity, we shall write

$$U_k := U_{\varepsilon_k, y_k}.$$

Estimate for $|U_k|_2^2$: For $N \geq 5$, $\frac{N}{N-2} < 2$. From (41),

$$|U_k|_2^2 = O({\varepsilon_k}^2). \tag{42}$$

Estimate for $|U_k|_{2^\#}^{2^\#}$: Since $y_k \in \partial\Omega$ and we are supposing that the domain is smooth,

$$|U_k|_{2^\#}^{2^\#} = \frac{2^\# B(N) \varepsilon_k}{2} + o(\varepsilon_k), \tag{43}$$

with

$$\begin{aligned} B(N) &= \frac{1}{2^\#} \int_{\mathbb{R}^N} U^{2^\#} \\ &= \omega_N \frac{1}{2^N} \sqrt{\pi} \frac{\Gamma\left(\frac{N}{2}\right)}{\Gamma\left(\frac{N+1}{2}\right)} [N(N-2)]^{\frac{N}{2}}, \end{aligned}$$

as proved in Appendix B. Here ω_N is the volume of the $N-1$ dimensional unit sphere.

Estimate for $|\nabla U_k|_2^2$ and for $|U_k|_{2^*}^{2^*}$: From Adimurthi and Mancini [1], since $N \geq 5$,

$$|\nabla U_k|_2^2 = \frac{S^{\frac{N}{2}}}{2} - \bar{C}_1 \varepsilon_k + O({\varepsilon_k}^2) \tag{44}$$

and

$$|U_k|_{2^*}^{2^*} = \frac{S^{\frac{N}{2}}}{2} - \bar{C}_2 \varepsilon_k + O({\varepsilon_k}^2), \tag{45}$$

where

$$\bar{C}_1 = H(y_k) \frac{\omega_{N-1}(N-2)^2}{4} \frac{\Gamma\left(\frac{N+3}{2}\right) \Gamma\left(\frac{N-3}{2}\right)}{\Gamma(N)} [N(N-2)]^{\frac{N-2}{2}}$$

and

$$\bar{C}_2 = H(y_k) \frac{\omega_{N-1}}{4} \frac{\Gamma\left(\frac{N+1}{2}\right) \Gamma\left(\frac{N-1}{2}\right)}{\Gamma(N)} [N(N-2)]^{\frac{N}{2}}.$$

Here $H(y_k)$ denotes the mean curvature of $\partial\Omega$ at y_k with respect to the unit outward normal and, as above, ω_N is the volume of the $N-1$ dimensional unit sphere. This yields

$$\frac{|\nabla U_k|_2^2}{|U_k|_{2^*}^2} = \frac{S}{2^{\frac{2}{N}}} - 2^{\frac{N-2}{N}} SH(y_k)A(N)\varepsilon_k + O(\varepsilon_k^2) \tag{46}$$

with

$$A(N) = \frac{2}{N}\frac{\omega_{N-1}}{\omega_N}\frac{\Gamma\left(\frac{N+1}{2}\right)\Gamma\left(\frac{N-3}{2}\right)}{\Gamma\left(\frac{N}{2}\right)\Gamma\left(\frac{N-2}{2}\right)} = \frac{N-1}{N}\frac{1}{\sqrt{\pi}}\frac{\Gamma\left(\frac{N-3}{2}\right)}{\Gamma\left(\frac{N-2}{2}\right)}. \tag{47}$$

To justify the last equality we recall that if $\omega_N(r)$ is the volume of the $N-1$ dimensional sphere with radius r, then

$$\omega_N(r) = \int_0^\pi \omega_{N-1}(r\sin\varphi)\, r d\varphi = r^{N-1}\omega_{N-1}(1)\int_0^\pi \sin^{N-2}\varphi\, d\varphi,$$

which yields

$$\frac{\omega_{N-1}}{\omega_N} = \frac{\omega_{N-1}(1)}{\omega_N(1)} = \frac{1}{\sqrt{\pi}}\frac{\Gamma\left(\frac{N}{2}\right)}{\Gamma\left(\frac{N-1}{2}\right)}.$$

We mention that the Talenti instanton we use does not coincide with the one in [1]. Denoting the Talenti instanton in Adimurthi and Mancini by V, $V(\,\cdot\,) = U((N(N-2))^{1/2}\,\cdot\,)$.

Estimate for $\int U_k w_k$:

Lemma 4.1.

$$\left|\int U_k w_k\right| \le \begin{cases} O\left(\varepsilon_k^{\frac{3}{2}}||w_k||\right) & \text{if } N=5, \\ O\left(\varepsilon_k^2|\log\varepsilon_k|^{\frac{2}{3}}||w_k||\right) & \text{if } N=6, \\ O\left(\varepsilon_k^2||w_k||\right) & \text{if } N\ge 7. \end{cases} \tag{48}$$

Proof.

$$\left|\int U_k w_k\right| \le |w_k|_{2^*}\left(\int U_k^{\frac{2N}{N+2}}\right)^{\frac{N+2}{2N}}.$$

If $N=5$, then $\frac{2N}{N+2} < \frac{N}{N-2}$. By (39),

$$\left|\int U_k w_k\right| \le C||w_k||\varepsilon_k^{\frac{N-2}{2}} = O\left(\varepsilon_k^{\frac{3}{2}}||w_k||\right).$$

If $N=6$, then $\frac{2N}{N+2} = \frac{N}{N-2}$. By (40),

$$\left|\int U_k w_k\right| \le C||w_k||\left(\varepsilon_k^{\frac{N}{2}}|\log\varepsilon_k|\right)^{\frac{N+2}{2N}} = O\left(\varepsilon_k^2|\log\varepsilon_k|^{\frac{2}{3}}||w_k||\right).$$

If $N\ge 7$, then $\frac{N}{N-2} < \frac{2N}{N+2}$. By (41),

$$\left|\int U_k w_k\right| \le C||w_k||\left(\varepsilon_k^{N\left(1-\frac{2N}{N+2}\frac{N-2}{2N}\right)}\right)^{\frac{N+2}{2N}} = O\left(\varepsilon_k^2||w_k||\right).$$

□

Estimate for $\int U_k^{2^*-1}w_k$*:* From [APY], Equations (3.15), for $N \geq 5$,

$$\int U_k^{2^*-1}w_k = O(\varepsilon_k||w_k||). \tag{49}$$

Estimate for $\int U_k^{2^\#-1}|w_k|$*:* Since $\frac{2N}{N+2} > 1$,

$$\begin{aligned}\int U_k^{2^\#-1}|w_k| &\leq |w_k|_{2^*}\left(\int U_k^{\frac{N}{N-2}\frac{2N}{N+2}}\right)^{\frac{N+2}{2N}} \\ &\leq C|w_k|_{2^*}\varepsilon_k^{\left(1-\frac{N}{N+2}\right)\frac{N+2}{2}} \\ &= O(\varepsilon_k||w_k||).\end{aligned} \tag{50}$$

Estimate for $\int U_k^{2^*-2}w_k^2$*:*

$$\int U_k^{2^*-2}w_k^2 = O(||w_k||^2). \tag{51}$$

Now we will obtain a lower bound for $I_{\alpha_k}(u_k)$. Let $v_k = u_k/C_k = U_k + \tilde{w}_k = U_k + w_k/C_k$. Because of (32), the sequence (v_k) satisfies (26) and the sequence $\tilde{w}_k$ satisfies (35). Of course, $d(v_k, M)$ is achieved by U_k. Because I is homogeneous of degree zero, $I_{\alpha_k}(u_k) = I_{\alpha_k}(v_k)$. We will compute $I_{\alpha_k}(v_k)$ but we will still call v_k by u_k, and $\tilde{w}_k$ by w_k.

Going back to (38), $I_\alpha(u_k)$ is bounded below by the sum of $\beta(u_k)$ and $\frac{4}{2^\#}\beta(u_k)\delta(u_k)$. We start by obtaining lower bounds for $\beta(u_k)$ and $\frac{4}{2^\#}\beta(u_k)\delta(u_k)$ separately. The expression for $\beta(u_k)$ involves two terms: $||u_k||^2$ and $|u_k|_{2^*}^{2^*}$. The first one is obviously

$$\begin{aligned}||u_k||^2 &= ||U_k||^2 + 2(\textstyle\int \nabla U_k \cdot \nabla w_k + a\int U_k w_k) + \ ||w_k||^2 \\ &= A_1 + A_2 + A_3.\end{aligned} \tag{52}$$

For the second term we use

Lemma 4.2 ([APY] Lemma 3.5)**.** *Let $q > 1$ and L be a non negative integer with $L \leq q$. Let V and ω be measurable functions on Ω with $V \geq 0$ and $V + \omega \geq 0$. Then*

$$\begin{aligned}\int (V+\omega)^q &= \sum_{i=0}^{L}\frac{q(q-1)\dots(q-i+1)}{i!}\int V^{q-i}\omega^i \\ &\quad + O\left(\int [V^{q-r}|\omega|^r + |\omega|^q]\right),\end{aligned}$$

where $r = \min\{L+1, q\}$.

Taking $L = 2$ and $q = 2^*$,

$$|u_k|_{2^*}^{2^*} = |U_k|_{2^*}^{2^*} + 2^*\int U_k^{2^*-1}w_k + \frac{2^*(2^*-1)}{2}\int U_k^{2^*-2}w_k^2 + O(||w_k||^r), \tag{53}$$

where $r = \min\{2^*, 3\}$, i.e., $r = 3$ if $N = 5$, and $r = 2^*$ if $N > 5$. The inequality

$$(1+z)^{-\eta} \geq 1 - \eta z, \tag{54}$$

for $\eta > 0$ and $z \geq -1$, implies

$$\begin{aligned} |u_k|_{2^*}^{-2} &\geq |U_k|_{2^*}^{-2}\left(1 - \frac{2\int U_k^{2^*-1}w_k}{|U_k|_{2^*}^{2^*}} \right. \\ &\qquad \left. - \frac{(2^*-1)\int U_k^{2^*-2}w_k^2}{|U_k|_{2^*}^{2^*}} + O(||w_k||^r)\right) \qquad (55) \\ &= B_1 + B_2 + B_3 + B_4. \end{aligned}$$

Let

$$l := \frac{|\nabla U_k|_2^2}{|U_k|_{2^*}^{2^*}}.$$

From (44) and (45),

$$l = 1 + O(\varepsilon_k). \qquad (56)$$

Using (52) and (55), we can write,

$$\beta(u_k) \geq \bar{I}_1 + \bar{I}_2 + \bar{I}_3 + \bar{I}_4,$$

where

$$\begin{aligned} \bar{I}_1 &= \frac{||U_k||^2}{|U_k|_{2^*}^2} \\ &= A_1 B_1 \\ \bar{I}_2 &= \frac{2}{|U_k|_{2^*}^2}\left[\int \nabla U_k \cdot \nabla w_k + a\int U_k w_k - l\int U_k^{2^*-1}w_k\right] \\ &= A_2 B_1 + A_1 B_2 \\ \bar{I}_3 &= \frac{1}{|U_k|_{2^*}^2}\left[||w_k||^2 - l(2^*-1)\int U_k^{2^*-2}w_k^2\right] \\ &= A_3 B_1 + A_1 B_3 \end{aligned}$$

and

$$\begin{aligned} \bar{I}_4 &= [(A_1 + A_3)B_4] + [A_2(B_2 + B_3 + B_4)] + A_3 B_2 + A_3 B_3 \\ &= E_1 + E_2 + E_3 + E_4. \end{aligned}$$

By (42) and (45),

$$\bar{I}_1 = \frac{|\nabla U_k|_2^2}{|U_k|_{2^*}^2} + o(\varepsilon_k).$$

We recall (35), $w_k \to 0$ in $H^1(\Omega)$.

By (34), the first of the four terms in $\bar{I}_2$ is zero; by Lemma 4.1 and by (49) the second and the third ones are $o(\varepsilon_k)$:

$$\bar{I}_2 = o(\varepsilon_k).$$

By (45), (51) and (56),

$$\bar{I}_3 = 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}}\left[||w_k||^2 - (2^*-1)\int U_k^{2^*-2}w_k^2\right]$$

The term E_1 is $o(||w_k||^2)$ because B_4 is $o(||w_k||^2)$. The term E_2 is $o(\varepsilon_k)$ because, from Lemma 4.1, A_2 is $o(\varepsilon_k)$. The term E_3 is $o(\varepsilon_k)$ because, from

(49), B_2 is $o(\varepsilon_k)$. Finally, the term E_4 is $o(||w_k||^2)$ because both A_3 and B_3 are $O(||w_k||^2)$. Therefore,

$$\bar{I}_4 = o(\varepsilon_k) + o(||w_k||^2).$$

Combining the expressions for $\bar{I}_1$, $\bar{I}_2$, $\bar{I}_3$ and $\bar{I}_4$,

$$\begin{aligned}\beta(u_k) &= \frac{|\nabla U_k|_2^2}{|U_k|_{2^*}^2} + 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} \left[||w_k||^2 - (2^*-1) \int U_k^{2^*-2} w_k^2 \right] \\ &\quad + o(\varepsilon_k) + o\left(||w_k||^2\right) \\ &\geq \frac{|\nabla U_k|_2^2}{|U_k|_{2^*}^2} + 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} \left[\gamma_2 ||w_k||^2 - (2^*-1) \int U_k^{2^*-2} w_k^2 \right] + o(\varepsilon_k),\end{aligned}$$

for any fixed number $\gamma_2 < 1$, because $a > 0$. This is our lower bound for $\beta(u_k)$.

Now we turn to the term $\frac{4}{2^\#}\beta(u_k)\delta(u_k)$ and write

$$\frac{4}{2^\#}\beta(u_k)\delta(u_k) = \frac{2}{2^\#} \frac{||u_k||}{|u_k|_{2^*}^{2+2^*/2}} \alpha_k |u_k|_{2^\#}^{2^\#} \tag{57}$$

We obtain a lower bound for $||u_k||$ from (52). Using (34), (42), (44) and Lemma 4.1,

$$||u_k|| \geq \left(\frac{S^{\frac{N}{2}}}{2} \right)^{\frac{1}{2}} + O(\varepsilon_k) + O(||w_k||^2).$$

We obtain a lower bound for $|u_k|_{2^*}^{-(2+2^*/2)}$ from (53). Using (45), (49), (51) and (54),

$$|u_k|_{2^*}^{-(2+2^*/2)} \geq \left(\frac{S^{\frac{N}{2}}}{2} \right)^{-\frac{1}{2}-\frac{2}{2^*}} + O(\varepsilon_k) + O(||w_k||^2).$$

For the product we obtain the lower bound

$$\begin{aligned}\frac{||u_k||}{|u_k|_{2^*}^{2+2^*/2}} &\geq 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} + O(\varepsilon_k) + O(||w_k||^2) \qquad (58) \\ &= D_1 + D_2 + D_3.\end{aligned}$$

To estimate the term $\alpha_k |u_k|_{2^\#}^{2^\#}$ we do not use Lemma 4.2 because it would give rise to a term $O\left(\alpha_k ||w_k||^{2^\#}\right)$, for which we do not have estimates. Instead we use this calculus

Lemma 4.3. *Let $\eta > 2$. For any $z \geq -1$,*

$$\frac{\eta(\eta-1)}{2} z^2 - \tilde{C}|z| + 1 \leq (z+1)^\eta, \tag{59}$$

where $\tilde{C} = 1 + \eta(\eta-1)/2$.

Proof. The difference between the right hand side and the left hand side is zero for $z = -1$ and $z = 0$. It is increasing for $z > 0$ and concave for $-1 < z < 0$. □

((59) also hold for $\eta = 2$, with equality for negative values of z.)

As a consequence of Lemma 4.3,

$$|u_k|_{2^\#}^{2^\#} \geq |U_k|_{2^\#}^{2^\#} - 2^\# \hat{C} \int U_k^{2^\#-1}|w_k| + \frac{2^\#(2^\#-1)}{2} \int U_k^{2^\#-2} w_k^2,$$

with

$$\hat{C} := \frac{\tilde{C}}{2^\#} = \frac{1}{2^\#} + \frac{2^\#-1}{2}.$$

Using (43) and (50),

$$\begin{aligned} \frac{2}{2^\#}\alpha_k |u_k|_{2^\#}^{2^\#} &\geq B(N)\alpha_k\varepsilon_k + (2^\#-1)\alpha_k \int U_k^{2^\#-2} w_k^2 + o(\alpha_k\varepsilon_k) \quad (60) \\ &= F_1 + F_2 + F_3. \end{aligned}$$

We will now substitute (58) and (60) in (57). On the one hand,

$$(D_1 + D_2 + D_3)F_3 = o(\alpha_k\varepsilon_k)$$

and

$$(D_2 + D_3)F_1 = o(\alpha_k\varepsilon_k).$$

On the other hand, by (36),

$$D_2 F_2 = O\left(\alpha_k \varepsilon_k^2 |\log \varepsilon_k|^{\frac{2}{N}} ||w_k||^2\right) = o(\alpha_k\varepsilon_k).$$

So,

$$\begin{aligned} \frac{4}{2^\#}\beta(u_k)\delta(u_k) &\geq 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} B(N)\alpha_k\varepsilon_k \\ &\quad + 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} (2^\#-1)\alpha_k \int U_k^{2^\#-2} w_k^2 \\ &\quad + O\left(||w_k||^2\right)\alpha_k \int U_k^{2^\#-2} w_k^2 + o(\alpha_k\varepsilon_k) \\ &\geq 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} \left[B(N)\alpha_k\varepsilon_k + \gamma_2(2^\#-1)\alpha_k \int U_k^{2^\#-2} w_k^2\right] \\ &\quad + o(\alpha_k\varepsilon_k), \end{aligned}$$

for any fixed number $\gamma_2 < 1$. This is our lower bound for $\frac{4}{2^\#}\beta(u_k)\delta(u_k)$.

Combining the lower bounds for $\beta(u_k)$ and for $\frac{4}{2^\#}\beta(u_k)\delta(u_k)$,

$$\begin{aligned} I_{\alpha_k}(u_k) &\geq \frac{|\nabla U_k|_2^2}{|U_k|_{2^*}^2} + 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} B(N)\alpha_k\varepsilon_k \\ &\quad + 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} \left[\gamma_2 ||w_k||^2 + \gamma_2(2^\#-1)\alpha_k \int U_k^{2^\#-2} w_k^2 \right. \\ &\quad \left. - (2^*-1)\int U_k^{2^*-2} w_k^2\right] + o(\alpha_k\varepsilon_k). \end{aligned}$$

From Lemma 3.11, the term inside the square parenthesis is greater than

$$\left[\left(\gamma_2 - \frac{(2^*-1)}{(2^*-1)+\gamma_1}\right)\left(||w_k||^2 + (2^\#-1)\alpha_k \int U_k^{2^\#-2} w_k^2\right) + o(\varepsilon_k)\right].$$

Choosing $\gamma_2 \geq \frac{(2^*-1)}{(2^*-1)+\gamma_1}$, yields that this term is greater than $o(\varepsilon_k)$. Hence,

$$I_{\alpha_k}(u_k) \geq \frac{|\nabla U_k|_2^2}{|U_k|_{2^*}^2} + 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} B(N)\alpha_k\varepsilon_k + o(\alpha_k\varepsilon_k).$$

Substituting (46) into this expression, we obtain

$$\begin{aligned} I_{\alpha_k}(u_k) &\geq \frac{S}{2^{\frac{2}{N}}} + 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} B(N)\alpha_k\varepsilon_k \left[1 - S^{\frac{N}{2}}\frac{A(N)}{B(N)}H(y_k)\frac{1}{\alpha_k} + o(1)\right] \\ &> \frac{S}{2^{\frac{2}{N}}}, \end{aligned}$$

for large k.

So assume α_0, in (18), is $+\infty$. Choose a sequence $\alpha_k \to +\infty$ as $k \to +\infty$ and denote by u_k a minimizer for I_{α_k} satisfying (1_{α_k}). From Lemmas 3.5 and 3.6, the conditions (26), (31), (32), (33) and (35) hold. Therefore $S_{\alpha_k} = I_{\alpha_k}(u_k) > \frac{S}{2^{\frac{2}{N}}}$ for large k, which is impossible. By Corollary 3.4, this establishes (ii) of Theorem 2.1.

Remark 4.4. *Since* $S^{\frac{N}{2}} = \int_{\mathbb{R}^N} U^{2^*} = \omega_N \frac{1}{2^N}\sqrt{\pi}\frac{\Gamma\left(\frac{N}{2}\right)}{\Gamma\left(\frac{N+1}{2}\right)}[N(N-2)]^{\frac{N}{2}}$, *it follows that*

$$B(N) = S^{\frac{N}{2}}.$$

Using

$$\omega_N = \frac{2\pi^{\frac{N}{2}}}{\Gamma\left(\frac{N}{2}\right)},$$

the common value is

$$B(N) = S^{\frac{N}{2}} = \frac{\pi^{\frac{N+1}{2}}}{2^{N-1}}\frac{1}{\Gamma\left(\frac{N+1}{2}\right)}[N(N-2)]^{\frac{N}{2}}.$$

5. Least energy solutions of (1_{α_0})

In this section we give a lower bound for $\alpha_0 = \min\left\{\alpha \mid S_\alpha = S/2^{\frac{2}{N}}\right\}$, and give partial results concerning existence of least energy solutions of (1_{α_0}).

From (10) we obtain

Lemma 5.1. *There exists a constant* $\bar{c} > \frac{4}{(2^\#)^{2/N}}$ *such that*

$$I_\alpha \leq \beta\left(1 + \frac{4}{2^\#}\delta + \bar{c}\delta^2\right). \tag{61}$$

Proof. Consider $\Lambda : [0, +\infty[\to \mathbb{R}$, defined by

$$\Lambda(\underline{\delta}) := \frac{1}{(2^\#)^{\frac{2}{N}}}\left[\left(\underline{\delta} + \sqrt{\underline{\delta}^2+1}\right)^N + \frac{2^*}{2}\left(\underline{\delta} + \sqrt{\underline{\delta}^2+1}\right)^{N-2}\right]^{\frac{2}{N}}.$$

Since $\frac{\partial}{\partial\underline{\delta}}\sqrt{\underline{\delta}^2+1}\Big|_{\underline{\delta}=0} = 0$ and $\frac{\partial}{\partial\underline{\delta}}\frac{1}{\sqrt{\underline{\delta}^2+1}}\Big|_{\underline{\delta}=0} = 0$, the first two derivatives of Λ at zero are

$$\Lambda'(0) = \frac{1}{(2^\#)^{\frac{2}{N}}}\frac{2}{N}(2^\#)^{\frac{2}{N}-1}\left[N + \frac{2^*}{2}(N-2)\right] = \frac{4}{2^\#}$$

and

$$\Lambda''(0) = \frac{1}{(2^\#)^{\frac{2}{N}}}\frac{2}{N}\left(\frac{2}{N}-1\right)(2^\#)^{\frac{2}{N}-2}(2N)^2 + \frac{2}{2^\#}[N + (N-2)] = \frac{4}{2^\#}\frac{2N-3}{N-1}.$$

Fix any number $c_1 > \frac{2}{2^\#}\frac{2N-3}{N-1}$. There exists an $\epsilon > 0$ such that (61) holds for $\bar{c} = c_1$ and $0 \le \underline{\delta} < \epsilon$.

Fix any number $c_2 > \frac{4}{(2^\#)^{2/N}}$. From (10), there exists an $L > 0$ such that (61) holds for $\bar{c} = c_2$ and $\underline{\delta} > L$.

The inequalities $\frac{2}{2^\#}\frac{2N-3}{N-1} < \frac{4}{2^\#} < \frac{4}{(2^\#)^{2/N}}$ show that $\max\left\{\frac{2}{2^\#}\frac{2N-3}{N-1}, \frac{4}{(2^\#)^{2/N}}\right\} = \frac{4}{(2^\#)^{2/N}}$.

By taking $\bar{c} \ge \max\{c_1, c_2\}$, $\bar{c}$ sufficiently large, we can guarantee (61) for all $\underline{\delta} \in [\epsilon, L]$. □

Lemma 5.2. *If $\alpha < A(N)\max_{\partial\Omega} H$, then $S_\alpha < \frac{S}{2^{\frac{2}{N}}}$.*

Proof. Choose $P \in \partial\Omega$ such that $H(P) = \max_{\partial\Omega} H$. From (42) and (46),

$$\beta(U_{\varepsilon,P}) = \frac{S}{2^{\frac{2}{N}}} - 2^{\frac{N-2}{N}} SH(P)A(N)\varepsilon + o(\varepsilon),$$

whereas, from (7) and (42)-(45),

$$\delta(U_{\varepsilon,P}) = \frac{2}{S^{\frac{N}{2}}}\frac{2^\#}{4}B(N)\alpha\varepsilon + o(\varepsilon).$$

The previous lemma implies that

$$\begin{aligned} S_\alpha &\le I_\alpha(U_{\varepsilon,P}) \\ &\le \frac{S}{2^{\frac{2}{N}}} - 2^{\frac{N-2}{N}} S^{\frac{2-N}{2}} B(N)\alpha\varepsilon\left[S^{\frac{N}{2}}\frac{A(N)}{B(N)}H(P)\frac{1}{\alpha} - 1 + o(1)\right] \\ &= \frac{S}{2^{\frac{2}{N}}} - 2^{\frac{N-2}{N}} S\alpha\varepsilon\left[A(N)H(P)\frac{1}{\alpha} - 1 + o(1)\right] \end{aligned}$$

as $\varepsilon \to 0$. Since, by assumption, $\alpha < A(N)\max_{\partial\Omega} H = A(N)H(P)$, $S_\alpha < \frac{S}{2^{\frac{2}{N}}}$. □

Corollary 5.3. *The value α_0 is greater than or equal to $A(N)\max_{\partial\Omega} H$.*

We let $|\Omega|$ denote the Lebesgue measure of Ω. By testing I_α with constant functions we obtain

Lemma 5.4. *If $a \le \frac{S}{(2|\Omega|)^{\frac{2}{N}}}$, then $\alpha_0 \ge \max\left\{\alpha \in [0, +\infty[\,\middle|\, I_\alpha(1) \le \frac{S}{2^{\frac{2}{N}}}\right\}$.*

Note. The value of $I_\alpha(1)$ is

$$I_\alpha(1) = \frac{|\Omega|^{\frac{2}{N}}}{(2^\#)^{\frac{2}{N}}}\left[\left(\frac{\alpha+\sqrt{\alpha^2+4a}}{2}\right)^N + \frac{2^*}{2}a\left(\frac{\alpha+\sqrt{\alpha^2+4a}}{2}\right)^{N-2}\right]^{\frac{2}{N}}.$$

We have not determined the exact value of α_0. However, using the ideas of Chabrowski and Willem [10], we have the following proposition concerning existence of least energy solutions for $\alpha = \alpha_0$:

Proposition 5.5. *If $\alpha_0 > A(N)\max_{\partial\Omega} H$ then there exists a least energy solution of (1_{α_0}).*

Proof. Choose a sequence $\alpha_k \nearrow \alpha_0$. Let u_k be a minimizer of I_{α_k} satisfying (1_{α_k}). As in the proof of Lemma 3.5, we conclude that the sequence (u_k) is bounded in $H^1(\Omega)$. So we can assume $u_k \rightharpoonup u$.

We claim that $u \neq 0$. Suppose, by contradiction, that $u = 0$. If the norms $|u_k|_{L^\infty(\Omega)}$ are uniformly bounded, then, from (26), $|u|_{2^*}^{2^*} = \frac{S^{\frac{N}{2}}}{2}$, which contradicts $u = 0$. If $|u_k|_{L^\infty(\Omega)} \to +\infty$, then Lemma 3.6 implies that we can repeat the argument of the previous sections to conclude that $S_{\alpha_k} > \frac{S}{2^{\frac{2}{N}}}$, for large k. This is also a contradiction. So $u \neq 0$.

Since $u \neq 0$, the argument in the proof of Lemmas 3.1 and 3.2 yields that u is a least energy solution of (1_{α_0}). Indeed, with the notations in the proof of Lemma 3.1, $x_0 \neq 0$. If $[h(1)]^{\frac{2}{N}} / [4(2^\#)^{\frac{2}{N}}] > \frac{S}{2^{\frac{2}{N}}}$, then $S_{\alpha_0} = [h(x_0)]^{\frac{2}{N}} / [4(2^\#)^{\frac{2}{N}}] > \frac{S}{2^{\frac{2}{N}}}$. Hence $I_{\alpha_0}(u) = [h(1)]^{\frac{2}{N}} / [4(2^\#)^{\frac{2}{N}}] = \frac{S}{2^{\frac{2}{N}}}$. □

Remark 5.6. *If a is sufficiently small and Ω is a (unit) ball, then the lower bound for α_0 in* Corollary 5.3 *is smaller than the lower bound for α_0 in* Lemma 5.4 *so that the previous proposition applies.*

Proof. The lower bound for α_0 in Corollary 5.3 is $A(N)$, given in (47). As $a \to 0$, the lower bound for α_0 in Lemma 5.4 tends to

$$\begin{aligned}
\left(\frac{(2^\#)^{\frac{2}{N}}}{|\Omega|^{\frac{2}{N}}} \frac{S}{2^{\frac{2}{N}}}\right)^{\frac{1}{2}} &= \left(\frac{2^\#}{2}\right)^{\frac{1}{N}} S^{\frac{1}{2}} \frac{1}{|\Omega|^{\frac{1}{N}}} \\
&= \left(\tfrac{N-1}{N-2}\right)^{\frac{1}{N}} \frac{\pi^{\frac{N+1}{2N}}}{2^{\frac{N-1}{N}}} \frac{1}{\left[\Gamma\left(\frac{N+1}{2}\right)\right]^{\frac{1}{N}}} [N(N-2)]^{\frac{1}{2}} \frac{\left[\Gamma\left(\frac{N+2}{2}\right)\right]^{\frac{1}{N}}}{\pi^{\frac{1}{2}}} \\
&= \left[\frac{\pi^{\frac{1}{2}}}{2^{N-1}} \left(\tfrac{N-1}{N-2}\right) \frac{\Gamma\left(\frac{N+2}{2}\right)}{\Gamma\left(\frac{N+1}{2}\right)}\right]^{\frac{1}{N}} [N(N-2)]^{\frac{1}{2}}.
\end{aligned}$$

□

Suppose now $\alpha_0 = A(N) \max_{\partial\Omega} H$. Once again, choose a sequence $\alpha_k \nearrow \alpha_0$ and let u_k be a minimizer of I_{α_k} satisfying (1_{α_k}). The argument in the proof of the previous proposition shows that, modulo a subsequence, either $u_k \rightharpoonup u \neq 0$, or $u_k \rightharpoonup 0$ and $|u_k|_{L^\infty(\Omega)} \to +\infty$. *We have not determined which of these alternatives holds.* In the first case u is a least energy solution of (1_{α_0}). In the second case let, as before, P_k be such that $u_k(P_k) = |u_k|_{L^\infty(\Omega)}$. Any limit point of (P_k) is contained in the set of points of maximum mean curvature of $\partial\Omega$. For if y_0 is a limit point of P_k, then

$$\begin{aligned}
-2^{\frac{N-2}{N}} SH(y_k)A(N)\varepsilon_k &= \left[2^{\frac{N-2}{N}} SH(y_0)A(N)\varepsilon_k - 2^{\frac{N-2}{N}} SH(y_k)A(N)\varepsilon_k\right] \\
&\quad -2^{\frac{N-2}{N}} SH(y_0)A(N)\varepsilon_k \\
&= -2^{\frac{N-2}{N}} SH(y_0)A(N)\varepsilon_k + o(\varepsilon_k).
\end{aligned}$$

If $H(y_0) < \max_{\partial\Omega} H$, then the argument in the previous section shows that $S_{\alpha_k} > \frac{S}{2^{\frac{2}{N}}}$, for large k.

We summarize these observations in

Proposition 5.7. *Suppose $\alpha_0 = A(N) \max_{\partial\Omega} H$. Then*

(i) either there exists a least energy solution of (1_{α_0}),

(ii) or any sequence, u_k, of least energy solutions of (1_{α_k}), for $\alpha_k < \alpha_0$, $\alpha_k \to \alpha_0$, has a subsequence, u_k, $u_k \rightharpoonup 0$, $|u_k|_{L^\infty(\Omega)} \to +\infty$; the limit points of any sequence of maximums of u_k are contained in the set of points of maximum mean curvature of the boundary of Ω.

Appendix A. The functional restricted to the Nehari manifold

In this Appendix we start by checking, using standard arguments, that the Nehari set $\mathcal{N}$ is a manifold and a natural constraint for Φ_α (defined in (2)). We then derive the expressions (3) and (6) for Φ_α restricted to $\mathcal{N}$, we derive an expression for I_α (defined in (8)) equivalent to (9) and to (10), and we derive upper and lower bounds for I_α.

Consider the set

$$\mathcal{N} := \left\{ u \in H^1(\Omega) : \Phi'_\alpha(u)u = 0, u \neq 0 \right\},$$

where Φ_α is the C^2 functional defined in (2), and define $J_\alpha : H^1(\Omega) \to \mathbb{R}$ by

$$J_\alpha(u) := \Phi'_\alpha(u)u = ||u||^2 + \alpha|u|^{2^\#}_{2^\#} - |u|^{2^*}_{2^*}.$$

The set $\mathcal{N} = \left\{ u \in H^1(\Omega) : J_\alpha(u) = 0, u \neq 0 \right\}$, is a manifold (called the Nehari manifold). Indeed, if $u \in \mathcal{N}$, then $J'_\alpha(u) \neq 0$, because if $J_\alpha(u) = 0$ and $J'_\alpha(u)u = 0$, then

$$0 = 2^* J_\alpha(u) - J'_\alpha(u)u = (2^* - 2)||u||^2 + (2^* - 2^\#)\alpha|u|^{2^\#}_{2^\#}.$$

This yields $u = 0$. Furthermore, the Nehari manifold is a natural constraint for Φ_α, by which we mean that any critical point of $\Phi_\alpha|_{\mathcal{N}}$ is a critical point of Φ_α. In fact, suppose that $u \in \mathcal{N}$ is a critical point of $\Phi_\alpha|_{\mathcal{N}}$. Then there exists a $\lambda \in \mathbb{R}$ such that $\Phi'_\alpha(u) = \lambda J'_\alpha(u)$. Applying both sides to u, $0 = J_\alpha(u) = \Phi'_\alpha(u)u = \lambda J'_\alpha(u)u$. However, we just saw that $J'_\alpha(u)u \neq 0$ if $J_\alpha(u) = 0$ (and $u \neq 0$). It follows that $\lambda = 0$ and u is a critical point of Φ_α.

For any $u \in H^1(\Omega) \setminus \{0\}$ there exists a unique $t(u) > 0$ such that $t(u)u \in \mathcal{N}$, i.e. $\Phi'_\alpha(t(u)u)t(u)u = 0$. The value of $t(u)$ is the solution of

$$||u||^2 + \alpha|u|^{2^\#}_{2^\#}[t(u)]^{2^\#-2} - |u|^{2^*}_{2^*}[t(u)]^{2^*-2} = 0.$$

Since $2^\# - 2 = \frac{2}{N-2}$ is half of $2^* - 2$, the equation

$$a + bt^{2^\#-2} - ct^{2^*-2} = 0.$$

is quadratic in $t^{\frac{2}{N-2}}$. Define the functionals a, b and $c : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$ by

$$\begin{array}{rcl} a(u) & := & ||u||^2, \\ b(u) & := & \alpha|u|^{2^\#}_{2^\#} \quad = \quad b_\alpha(u), \\ c(u) & := & |u|^{2^*}_{2^*}. \end{array}$$

(Note that $a \neq a$.) The value of $t(u)$ is

$$t(u) = \left(\frac{b + \sqrt{b^2 + 4ac}}{2c} \right)^{\frac{N-2}{2}} (u). \tag{62}$$

The functional $t : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$ is obviously continuous and the map $u \mapsto t(u)u$ defines a homeomorphism of the unit sphere in $H^1(\Omega)$ with $\mathcal{N}$. Its inverse is the retraction $u \mapsto \frac{u}{||u||}$.

We define $\Psi_\alpha : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$ by

$$\Psi_\alpha(u) := \Phi_\alpha(t(u)u).$$

In terms of a, b, c and t,

$$\Psi_\alpha = \frac{1}{2}at^2 + \frac{1}{2^\#}bt^{2^\#} - \frac{1}{2^*}ct^{2^*}.$$

Replacing (62) into this expression for Ψ_α, and simplifying, leads to

$$\Psi_\alpha = \frac{1}{N}\frac{1}{2^\#}\left[\left(\frac{b+\sqrt{b^2+4ac}}{2c}\right)^N c + \frac{2^*}{2}\left(\frac{b+\sqrt{b^2+4ac}}{2c}\right)^{N-2} a\right].$$

We now introduce the functionals β, $\gamma : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$, defined by

$$\beta := \frac{a}{c^{\frac{N-2}{N}}}$$

and

$$\gamma = \gamma_\alpha := \frac{b}{c^{\frac{N-1}{N}}},$$

as in expressions (4) and (5), respectively. In terms of β and γ, the expression for Ψ_α is

$$\Psi_\alpha = \frac{1}{N}\frac{1}{2^\#}\frac{1}{2^N}\left[\left(\gamma+\sqrt{\gamma^2+4\beta}\right)^N + 2\cdot 2^*\beta\left(\gamma+\sqrt{\gamma^2+4\beta}\right)^{N-2}\right].$$

This is (3). If we introduce still another functional $\delta : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$, defined by

$$\delta = \delta_\alpha := \frac{\gamma}{2\sqrt{\beta}},$$

as in expression (7), then we can write Ψ_α as

$$\Psi_\alpha = \frac{1}{N}\frac{\beta^{\frac{N}{2}}}{2^\#}\left[\left(\delta+\sqrt{\delta^2+1}\right)^N + \frac{2^*}{2}\left(\delta+\sqrt{\delta^2+1}\right)^{N-2}\right].$$

This is (6).

We give an expression for $I_\alpha = (N\Psi_\alpha)^{\frac{2}{N}}$, defined in (8), equivalent to (9) and to (10):

$$I_\alpha = \beta\left(\delta+\sqrt{\delta^2+1}\right)^{\frac{4}{2^*}}\left(\frac{2}{2^\#}\delta^2 + \frac{2}{2^\#}\delta\sqrt{\delta^2+1}+1\right)^{\frac{2}{N}}.$$

Since

$$\frac{4}{2^*} + \frac{2}{N}\frac{2}{2^\#} = \frac{4}{2^\#},$$

I_α has the lower bound

$$I_\alpha \geq \beta\left(1+\frac{4}{2^\#}\delta\right). \tag{63}$$

For an upper bound for I_α we refer to Lemma 5.1.

Appendix B. The estimate for $|U_k|^{2^\#}_{2^\#}$

In this Appendix we use the ideas of Adimurthi and Mancini [1] to prove (43).

We wish to estimate $|U_{\varepsilon,y}|^{2^\#}_{2^\#}$, where $U_{\varepsilon,y}$ is defined in (14) and $y \in \partial\Omega$. By a change of coordinates we can assume that $y = 0$,

$$B_R(0) \cap \Omega = \{(x', x_N) \in B_R(0) | x_N > \rho(x')\}$$

and

$$B_R(0) \cap \partial\Omega = \{(x', x_N) \in B_R(0) | x_N = \rho(x')\},$$

for some $R > 0$, where $x' = (x_1, \ldots, x_{N-1})$,

$$\rho(x') = \sum_{i=1}^{N-1} \lambda_i x_i^2 + O(|x'|^3),$$

$\lambda_i \in \mathbb{R}$, $1 \le i \le N-1$.

We begin by supposing all the λ_i's are positive. Let $U_\varepsilon := U_{\varepsilon,0}$ and $\Sigma := \{(x', x_N) \in B_R(0) | 0 < x_N < \rho(x')\}$. Then

$$|U_\varepsilon|^{2^\#}_{2^\#} = \frac{1}{2}\int_{B_R(0)} U_\varepsilon^{2^\#} - \int_\Sigma U_\varepsilon^{2^\#} + \int_{B_R^C(0)\cap\Omega} U_\varepsilon^{2^\#}. \tag{64}$$

We will estimate each of the three terms on the right hand side of (64). For the third term we have

$$\begin{aligned}
\int_{B_R^C(0)\cap\Omega} U_\varepsilon^{2^\#} &\le \int_{B_R^C(0)} U_\varepsilon^{2^\#} \\
&= O\left(\varepsilon \int_{R/\varepsilon}^{+\infty} \frac{r^{N-1}}{(1+r^2)^{N-1}} dr\right) \\
&= O(\varepsilon \times \varepsilon^{N-2}) \\
&= O(\varepsilon^{N-1})
\end{aligned}$$

Using this estimate, for the first term on the right hand side of (64) we have

$$\begin{aligned}
\frac{1}{2}\int_{B_R(0)} U_\varepsilon^{2^\#} &= \frac{1}{2}\int_{\mathbb{R}^N} U_\varepsilon^{2^\#} + O(\varepsilon^{N-1}) \\
&= \frac{1}{2}\varepsilon \int_{\mathbb{R}^N} U^{2^\#} + O(\varepsilon^{N-1}) \\
&= \frac{2^\#}{2} B(N)\varepsilon + O(\varepsilon^{N-1}),
\end{aligned}$$

with

$$\begin{aligned} B(N) &:= \frac{1}{2^\#}\int_{\mathbb{R}^N} U^{2^\#} \\ &= \frac{1}{2^\#}\,\omega_N \int_0^{+\infty} \frac{r^{N-1}}{(1+r^2)^{N-1}}\,dr \times [N(N-2)]^{\frac{N}{2}} \\ &= \tfrac{N-2}{2(N-1)}\,\omega_N \times \frac{1}{2^{N-1}}\sqrt{\pi}\,\frac{\Gamma\left(\frac{N-2}{2}\right)}{\Gamma\left(\frac{N-1}{2}\right)} \times [N(N-2)]^{\frac{N}{2}} \\ &= \omega_N \frac{1}{2^N}\sqrt{\pi}\,\frac{\Gamma\left(\frac{N}{2}\right)}{\Gamma\left(\frac{N+1}{2}\right)}[N(N-2)]^{\frac{N}{2}}; \end{aligned}$$

here ω_N is the volume of the $N-1$ dimensional unit sphere.

So we are left with the estimate of the second term on the right hand side of (64). Let $\sigma > 0$ be such that

$$L_\sigma := \{x \in \mathbb{R}^N |\, |x_i| < \sigma, 1 \le i \le N\} \subset B_{\frac{R}{4}}(0)$$

and define

$$\Delta_\sigma := \{x' |\, |x_i| < \sigma, 1 \le i \le N-1\}.$$

For the second term on the right hand side of (64),

$$\begin{aligned} \int_\Sigma U_\varepsilon^{2^\#} &= \int_{\Sigma\cap L_\sigma} U_\varepsilon^{2^\#} + O(\varepsilon^{N-1}) \\ &= \int_{\Delta_\sigma}\int_0^{\rho(x')} U_\varepsilon^{2^\#}\,dx_N\,dx' + O(\varepsilon^{N-1}) \\ &= O\left(\int_{\Delta_\sigma}\int_0^{\rho(x')} \frac{\varepsilon^{N-1}}{(\varepsilon^2+|x|^2)^{N-1}}\,dx_N\,dx'\right) + O(\varepsilon^{N-1}); \end{aligned}$$

using the change of variables $\sqrt{\varepsilon^2+|x'|^2}\,y_N = x_N$,

$$\begin{aligned} &= O\left(\int_{\Delta_\sigma} \frac{\varepsilon^{N-1}}{(\varepsilon^2+|x'|^2)^{N-\frac{3}{2}}}\int_0^{\frac{\rho(x')}{\sqrt{\varepsilon^2+|x'|^2}}} \frac{1}{(1+y_N^2)^{N-1}}\,dy_N\,dx'\right) \\ &\quad + O(\varepsilon^{N-1}); \end{aligned}$$

since $\int_0^s \frac{1}{(1+t^2)^{N-1}}\,dt = s + O(s^3)$,

$$\begin{aligned} &= O\left(\varepsilon^{N-1}\int_{\Delta_\sigma} \frac{\sum \lambda_i x_i^2}{(\varepsilon^2+|x'|^2)^{N-1}}\,dx'\right) \\ &\quad + O\left(\varepsilon^{N-1}\int_{\Delta_\sigma} \frac{|x'|^3}{(\varepsilon^2+|x'|^2)^{N-1}}\,dx'\right) \\ &\quad + O(\varepsilon^{N-1}) \\ &= O\left(\varepsilon^2\int_{\Delta_\sigma/\varepsilon} \frac{|y'|^2}{(1+|y'|^2)^{N-1}}\,dy'\right) \\ &\quad + O\left(\varepsilon^3\int_{\Delta_\sigma/\varepsilon} \frac{|y'|^3}{(1+|y'|^2)^{N-1}}\,dy'\right) \\ &\quad + O(\varepsilon^{N-1}) \\ &= O(\varepsilon^2). \end{aligned}$$

Combining the estimates for the three terms on the right hand side of (64),

$$|U_\varepsilon|_{2^\#}^{2^\#} = \frac{2^\#}{2} B(N)\varepsilon + O(\varepsilon^2), \tag{65}$$

if all the λ_i's are positive. If all the λ_i's are negative, then the minus sign on the right hand side of (64) turns into a plus sign, and (65) follows. From these two cases we deduce that (65) holds no matter what the sign of the λ_i's is.

Acknowledgments. This work was initiated during the visit of the first author to Instituto Superior Técnico, whose kind hospitality and support he gratefully acknowledges.

References

[1] Adimurthi and G. Mancini, The Neumann problem for elliptic equations with critical nonlinearity, A tribute in honour of G. Prodi, *Scuola Norm. Sup. Pisa* (1991), 9-25.
[2] Adimurthi, F. Pacella and S.L. Yadava, Interaction between the geometry of the boundary and positive solutions of a semilinear Neumann problem with critical nonlinearity, *J. Funct. Anal.* **113** (1993), 318-350.
[3] Adimurthi and S.L. Yadava, Existence and Nonexistence of Positive Radial Solutions of Neumann Problems with Critical Sobolev Exponents, *Arch. Rational Mech. Anal.* **115** (1991), 275-296.
[4] Adimurthi and S.L. Yadava, Some remarks on Sobolev type inequalities, *Calc. Var.* **2** (1994), 427-442.
[5] S. Agmon, A. Douglis and L. Nirenberg, Estimates Near the Boundary for Solutions of Elliptic Partial Differential Equations Satisfying General Boundary Conditions. I, *Comm. Pure Appl. Math.* **12** (1959), 623-727.
[6] T. Aubin, "Nonlinear Analysis on Manifolds. Monge-Ampère equations," Grundlehren der Mathematischen Wissenschaften, Vol. 252, Springer-Verlag, New York, 1982.
[7] G. Bianchi and H. Egnell, A note on the Sobolev inequality, *J. Funct. Anal.* **100** (1991), 18-24.
[8] H. Brézis and E. Lieb, Sobolev inequalities with remainder terms, *J. Funct. Anal.* **62** (1985), 73-86.
[9] H. Brézis and L. Nirenberg, Positive Solutions of Nonlinear Elliptic Equations Involving Critical Sobolev Exponents, *Comm. on Pure and Appl. Math.* **36** (1983), 437-477.
[10] J. Chabrowski and M. Willem, Least energy solutions for the critical Neumann problem with a weight, to appear in *Calc. Var.*
[11] P. Cherrier, Problémes de Neumann nonlinéaires sur des variétés Riemannienes, *J. Funct. Anal.* **57** (1984), 154-207.
[12] B. Gidas and J. Spruck, A Priori bounds for positive solutions of nonlinear elliptic equations, *Comm. in PDE's* 6 No. **8** (1981), 883-901.
[13] P.M. Girão, A sharp inequality for Sobolev functions, *C. R. Acad. Sci. Paris, Sér. I* **334** (2002), 105-108.
[14] P.M. Girão, A family of sharp inequalities for Sobolev functions, to appear.
[15] P.L. Lions, The concentration-compactness principle in the calculus of variations, The limit case, *Revista Math. Iberoamericana* **1** No. 1 and No. 2 (1985), 145-201 and 45-120.
[16] O. Rey, The role of the Green's function in a nonlinear elliptic equation involving the critical Sobolev exponent, *J. Funct. Anal.* **89** (1990), 1-52.
[17] G. Talenti, Best constant in Sobolev Inequality, *Ann. Mat. Pura Appl.* **110** No. 4 (1976), 353-372.
[18] X.J. Wang, Neumann problems of semilinear elliptic equations involving critical Sobolev exponents, *J. Diff. Eq.* **93** (1991), 283-310.

[19] Z.Q. Wang, Existence and nonexistence of G-least energy solutions for a nonlinear Neumann problem with critical exponent in symmetric domains, *Calc. Var.* **8** (1999), 109-122.
[20] M. Zhu, Sharp Sobolev inequalities with interior norms, *Calc. Var.* **8** (1999), 27-43.
[21] M. Zhu, Uniqueness Results through a Priori Estimates I. A Three Dimensional Neumann Problem, *J. Differential Equations* **154** (1999), 284-317.

Department of Mathematical Sciences, University of Nevada at Las Vegas, Las Vegas, Nevada 89154-4020, USA
E-mail address: costa@nevada.edu

Mathematics Department, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisbon, Portugal
E-mail address: pgirao@math.ist.utl.pt